PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C12N 15/40, C07K 14/18, C12Q 1/70, C07K 16/10, G01N 33/576

(11) International Publication Number:

WO 96/13590

A2

(43) International Publication Date:

9 May 1996 (09.05.96)

(21) International Application Number:

PCT/EP95/04155

(22) International Filing Date:

23 October 1995 (23.10.95)

(30) Priority Data:

94870166.9 21 October 1994 (21.10.94) EP
(34) Countries for which the regional or
international application was filed: AT et al.
95870076.7 28 June 1995 (28.06.95) EP
(34) Countries for which the regional or

international application was filed:

AT et al.

(71) Applicant (for all designated States except US): INNOGENET-ICS N.V. [BE/BE]; Industriepark Zwijnaarde 7, Box 4, B-9052 Zwijnaarde (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MAERTENS, Geert [BE/BE]; Zilversparrenstraat 64, B-8310 Brugge (BE). STUYVER, Lieven [BE/BE]; Holestraat 8, B-2400 Mol (BE).

(74) Common Representative: INNOGENETICS N.V.; Mrs. Ann De Clercq, Industriepark Zwijnaarde 7, Box 4, B-9052 Zwijnaarde (BE). (81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).

Published

Without international search report and to be republished upon receipt of that report.

With a request for rectification under Rule 91.1(f).

(54) Title: NEW SEQUENCES OF HEPATITIS C VIRUS GENOTYPES AND THEIR USE AS PROPHYLACTIC, THERAPEUTIC AND DIAGNOSTIC AGENTS

(57) Abstract

The present invention relates to new genomic nucleotide sequences and amino acid sequences corresponding to the coding region of these genomes. The invention relates to new HCV types and subtypes sequences which are different from the known HCV types and subtypes sequences. More particularly, the present invention relates to new HCV type 7 sequences, new HCV type 9 sequences, new HCV type 10 and new HCV type 11 sequences. Also, the present invention relates to new HCV type 1 sequences of subtypes 1d, 1e, 1f and 1g; new HCV type 2 sequences of subtypes 2e, 2f, 2g, 2h, 2i, 2k and 2l; new HCV type 3 sequences of subtype 3g, new HCV type 4 sequences of subtypes 4k, 4l and 4m; a process for preparing them, and their use for diagnosis, prophylaxis and therapy. More particularly, the present invention provides new type-specific sequences of the Core, the E1 and the NS5 regions of new HCV types 7, 9, 10 and 11, as well as of new variants (subtypes) of HCV types 1, 2, 3 and 4. These new HCV sequences are useful to diagnose the presence of HCV type 1, and/or type 2, and/or type 3, and/or type 4, and/or type 9, and/or type 9, and/or type 10, and/or type 11 genotypes or serotypes in a biological sample. Moreover, the availability of these new type-specific sequences can increase the overall sensitivity of HCV detection and should also prove to be useful for prophylactic and therapeutic purposes.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy ·	PL	Poland
-		JP.	Japan	PT	Portugal
BR	Brazil ,·	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic	N.F	of Korea	SE	Sweden
CG	Congo	7/0	Republic of Korea	SI	Slovenia
CH	Switzerland	KR	Kazakhatan	SK	Slovakia
CI	Côte d'Ivoire	KZ		SN	
CM	Cameroon	L	Liechtenstein		Senegal
CN	China	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Latvin	T.J	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
es Pi	Finland	ML	Mali	UZ	Uzbekistan
		MN	Mongolia	VN	Viet Nam
FR	France	14814	1,101,94.00		
GA	Gabon				

10

15

20

25

30

NEW SEQUENCES OF HEPATITIS C VIRUS GENOTYPES AND THEIR USE AS PROPHYLACTIC, THERAPEUTIC AND DIAGNOSTIC AGENTS

The invention relates to new sequences of hepatitis C virus (HCV) genotypes and their use as prophylactic, therapeutic and diagnostic agents.

The present invention relates to new genomic nucleotide sequences and amino acid sequences corresponding to the coding region of these genomes. The invention relates to new HCV types and subtypes sequences which are different from the known HCV types and subtypes sequences. More particularly, the present invention relates to new HCV type 7 sequences, new HCV type 9 sequences, new HCV types 10 and new HCV type 11 sequences. Also the present invention relates to new HCV type 1 sequences of subtypes 1d, 1e, 1f and 1g; new HCV type 2 sequences of subtypes 2e, 2f, 2g, 2h, 2i, 2k and 2l; new HCV type 3 sequences of subtype 3g, new HCV type 4 sequences of subtypes 4k, 4l and 4m; a process for preparing them, and their use for diagnosis, prophylaxis and therapy.

The technical problem underlying the present invention is to provide new HCV sequences from untill now unknown HCV types and/or subtypes. More particularly, the present invention provides new type-specific sequences of the Core, the E1 and the NS5 regions of new HCV types 7, 9, 10 and 11, as well as of new variants (subtypes) of HCV types 1, 2, 3 and 4. These new HCV sequences are useful to diagnose the presence of HCV type 1, and/or type 2, and/or type 3, and/or type 4, and/or type 7, and/or type 9, and/or type 10, and/or type 11 genotypes or serotypes in a biological sample. Moreover, the availability of these new type-specific sequences can increase the overall sensitivity of HCV detection and should also prove to be useful for prophylactic and therapeutic purposes.

Hepatitis C viruses (HCV) have been found to be the major cause of non-A, non-B hepatitis. The sequences of cDNA clones covering the complete genome of several prototype isolates have been determined (Kato et al., 1990; Choo et al., 1991; Okamoto et al., 1992). Comparison of these isolates shows that the variability in nucleotide sequences can be used to distinguish at least 2 different genotypes, type 1 (HCV-1 and HCV-J) and type 2 (HC-J6 and HC-J8),

10

15

20

25

30

with an average homology of about 68%. Within each type, at least two subtypes exist (e.g. represented by HCV-1 and HCV-J), having an average homology of about 79%. HCV genomes belonging to the same subtype show average homologies of more than 90% (Okamoto et al., 1992). However, the partial nucleotide sequence of the NS5 region of the HCV-T isolates showed at most 67% homology with the previously published sequences, indicating the existence of yet another HCV type (Mori et al., 1992). Parts of the 5' untranslated region (UR), core, NS3, and NS5 regions of this type 3 have been published, further establishing the similar evolutionary distances between the 3 major genotypes and their subtypes (Chan et al., 1992). Type 4 was subsequently discovered (Stuyver et al., 1993b; Simmonds et al., 1993a; Bukh et al., 1993; Stuyver et al., 1994a). As well as type 5 (Stuyver et al., 1993b; Simmonds et al., 1993c; Bukh et al., 1993; Stuyver et al., 1994b), and type 6 HCV groups (Bukh et al., 1993; Simmonds et al., 1993c). An overview of the present state of the art regarding HCV genotypes is given in Table 3. The nomenclature system proposed by the inventors of the present application has now been accepted by scientists worldwide (Simmonds et al., 1994).

The aim of the present invention is to provide new HCV nucleotide and amino acid sequences enabling the detection of HCV infection.

Another aim of the present infection is to provide new nucleotide and amino acid HCV sequences enabling the classification of infected biological fluids into different serological groups.

Another aim of the present invention is to provide new nucleotide and amino acid HCV sequences ameliorating the overall HCV detection rate.

Another aim of the present invention is to provide new HCV sequences, useful for the design of HCV prophylactic or therapeutic vaccine compositions.

Another aim of the present invention is to provide a pharmaceutical composition consisting of antibodies raised against the polypeptides encoded by these new HCV sequences, for therapy or diagnosis.

All the aims of the present invention are met by the following embodiments of the present invention.

The present invention relates more particularly to an HCV polynucleic acid, having a nucleotide sequence which is unique to a heretofore unidentified HCV type or subtype which is different from HCV subtypes 1a, 1b, 1c, 2a, 2b, 2c, 2d, 3a, 3b,

10

15

20

25

30

3c, 3d, 3e, 3f, 4a, 4b, 4c, 4d, 4e, 4f, 4g, 4h, 4i, 4j, 5a or 6a, with said HCV subtypes being classified as in Table 3 by comparison of a part of the NS5 gene nucleotide sequence spanning positions 7932 to 8271, with said amino acid numbering being shown in Table 1, and with said polynucleic acid containing at least one nucleotide differing from said known HCV nucleotide sequences, or the complement thereof. The sequence of known HCV isolates may be found in any nucleotide sequence database known in the art (such as for instance the EMBL database).

The present invention thus also relates to a polynucleic acid having a nucleotide sequence which is unique to at least one of HCV subtypes 1d, 1e, 1f, 1g, 2e, 2f, 2g, 2h, 2i, 2k, 2l, 3g, 4k, 4l, 4m, 7a, 7c or 7d, with said HCV subtypes being classified as defined above.

The present invention thus also relates to a polynucleic acid having a nucleotide sequence which is unique to at least one of HCV types 9, 10 or 11, with said HCV types being classified as defined above.

It is to be noted that the nucleotide(s) difference in the polynucleic acids of the invention may involve an amino acid difference in the corresponding amino acid sequences encoded by said polynucleic acids. A composition according to the present invention may contain only polynucleic acid sequences or polynucleic acid sequences mixed with any excipient known in the art of diagnosis, prophylaxis or therapy.

According to a preferred embodiment, the present invention relates to a polynucleic acid encoding an HCV polyprotein comprising in its amino acid sequence at least one of the following amino acid residues:

I15, C38, V44, A49, Q43, P49, Q55, A58, S60 or D60, E68 or V68, H70, A71 or Q71 or N71, D72, H81, H101, D106, S110, L130, I134, E135, L140, S148, T150 or E150, Q153, F155, D157, G160, E165, I169, F181, L186, T190, T192 or I192 or H192, I193, A195, S196, R197 or N197 or K197, Q199 or D199 or H199 or N199, F200 or T200, A208, I213, M216 or S216, N217 or S217 or G217 or K217, T218, I219, A222, Y223, I230, W231 or L231, S232 or H232 or A232, Q233, E235 or L235, F236 or T236, F237, L240 or M240, A242, N244, N249, I250 or K250 or R250, A252 or C252, A254, I255 or V255, D256 or M256, E257, E260 or K260, R261, V268, S272 or R272, I285, G290 or F290, A291, A293 or L293

10

15

20

or W293, T294 or A294, S295 or H295, K296 or E296, Y297 or M297, I299 or Y299, I300, S301, P316, S2646, A2648, G2649, A2650, V2652, Q2653, H2656 or L2656, D2657, F2659, K2663 or Q2663, A2667 or V1667, D2677, L2681, M2686 or Q2686 or E2686, A2692 or K2692, H2697, I2707, L2708 or Y2708, A2709, A2719 or M2719, F2727, T2728 or D2728, E2729, F2730 or Y2730, I2741, I2745, V2746 or E2746 or L2746 or K2746, A2748, S2749 or P2749, R2750, E2751, D2752 or N2752 or S2752 or T2752 or V2752 or I2752 or Q2752, S2753 or D2753 or G2753, D2754, A2755, L2756 or Q2756, R2757,

with said notation being composed of a letter representing the amino acid residue by its one-letter code, and a number representing the amino acid numbering according to Kato et al. (1980), as shown in Table 1,

or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined in Table 5, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.

Each of the above-mentioned residues can be found in Figures 2, 4 or 6 showing the new amino acid sequences of the present invention aligned with known sequences of other types or subtypes of HCV for the Core/E1 region.

According to another preferred embodiment, the present invention relates to a polynucleic acid encoding a HCV polyprotein comprising in its amino acid sequence at least one amino acid sequence chosen from the following list:

ARQSDGRSWAQ or ARRSEGRSWAQ as for subtype 1d (SEQ ID NO 107 and 108)

	ERRPEGRSWAQ as for subtype 1e	(SEQ ID NO 109)
	ARRPEGRSWAQ as for subtype 1f	(SEQ ID NO 110)
25	DRRTTGKSWGR as for subtype 2k	(SEQ ID NO 111)
	DRRATGRSWGR as for subtype 2e	(SEQ ID NO 112)
	DRRATGKSWGR as for subtype 2f	(SEQ ID NO 113)
	VRQPTGRSWGQ as for type 9	(SEQ ID NO 114)
	VRHQTGRTWAQ as for subtype 7a and 7c	(SEQ ID NO 115)
30	VRQNQGRTWAQ as for subtype 7d	(SEQ ID NO 116)
	ARRTEGRSWAQ as for type 10	(SEQ ID NO 117)
	VRRTTGRXXXX or VRRTTGRTWAQ as for type 11	(SEQ ID NO 118
	and 119)	

	HEVRNASGVYHV or HEVRNASGVYHL as for subtype and 121)	1d (SEQ ID NO 120
	YEVHSTTDGYHV as for subtype 1f	(SEQ ID NO 122)
r	VEVKNTSQAYMA as for subtype 2e	(SEQ ID NO 123)
5	IQVKNNSHFYMA as for subtype 2f	(SEQ ID NO 124)
	VQVKNTSTMYMA as for subtype 2g	(SEQ ID NO 125)
	VQVKNTSHSYMV as for subtype 2h	(SEQ ID NO 126)
	VQVANRSGSYMV as for subtype 2i	(SEQ ID NO 127)
	VEIKNTXNTYVL or VEIKNTSNTYVL as for subtype 2k	(SEQ ID NO 128
10	and 129)	
	INYRNVSGIYYV or INYRNTSGIYHV or INYHNTSGIYHI o	r TNYRNVSGIYHV as
	for subtype 4k	(SEQ ID NO 130,
	131, 132 or 133)	
	QHYRNVSGIYHV as for subtype 4I	(SEQ ID NO 134)
15	IQVKNASGIYHL as for type 9	(SEQ ID NO 135)
	AHYTNKSGLYHL as for subtype 7c	(SEQ ID NO 136)
	LNYANKSGLYHL as for subtype 7d	(SEQ ID NO 137)
	LEYRNASGLYMV as for type 10	(SEQ ID NO 138)
	IYEMDGMIMHY or IYEMSGMILHA as for subtype 1d	(SEQ ID NO 139
20	and 140)	
	VYEAKDIILHT as for subtype 1f	(SEQ ID NO 141)
	VWQLXDAVLHV as for subtype 2e	(SEQ ID NO 142)
	VWQLRDAVLHV as for subtype 2f	(SEQ ID NO 143)
	IWQMQGAVLHV as for subtype 2g	(SEQ ID NO 144)
25	VWQLKDAVLHV as for subtype 2h	(SEQ ID NO 145)
	VWQLEEAVLHV as for subtype 2i	(SEQ ID NO 146)
	TWQLXXAVLHV as for subtype 2k	(SEQ ID NO 147)
	VYEADHHILHL or VYEADHHILAL or VFEADHHILHL as for	or subtupe 4k (SEQ
	ID NO 148, 149 and 150)	. ,
30	VYESDHHILHL as for subtype 4I	(SEQ ID NO
	151)	, , , , , , , , , , , , , , , , , , , ,
	VFEAETMILHL as for type 9	(SEQ ID NO 152)
	VYEAETLILHL as for subtype 7c	(SEQ ID NO
	• •	125612140

	153)	
	VYEANGMILHL as for subtype 7d	(SEQ ID NO 154)
	VYEAGDIILHL as for type 10	(SEQ ID NO 155)
	VREDNHLRCWMAL or VRENNSSRCWMAL as for sub	otype 1d
5	(SEQ ID	NO 156 and 157)
	IREGNISRCWVPL as for subtype 1f	(SEQ ID NO 158)
	ENSSGRFHCWIPI as for subtype 2e	(SEQ ID NO 159)
	ERSGNRTFCWTAV as for subtype 2f	(SEQ ID NO 160)
	ELQGNKSRCWIPV as for subtype 2g	(SEQ ID NO 162)
10	ERHQNQSRCWIPV as for subtype 2h	(SEQ ID NO 163)
	EWKDNTSRCWIPV as for subtype 2i	(SEQ ID NO 164)
	EREGNSSRCWIPV as for subtype 2k	(SEQ ID NO 165)
	VREGNOSRCWVAL or VRTGNOSRCWVAL or V	/RVGNQSSCWVAL or
	VRVGNQSRCWVAL or VKEGNHSRCWVAL as for sub	otype 4k
15	(SEQ ID NO 1	66, 167, 168 or 169)
	VKTGNTSRCWVAL as for subtype 4I	(SEQ ID NO 170)
	IKAGNESRCWLPV as for type 9	(SEQ ID NO 171)
	VKEGNQSRCWVQA as for subtype 7c	(SEQ ID NO 172)
	VKXXNLTKCWLSA as for subtype 7d	(SEQ ID NO 173)
20	VRSGNTSRCWIPV as for type 10	(SEQ ID NO 174)
	VKNASVPTAA or VKDANVPTAA as for subtype 1d	(SEQ ID NO 175
	and 176)	
	ARIANAPIDE as for subtype 1f	(SEQ ID NO 177)
	VSKPGALTKG as for subtype 2e	(SEQ ID NO 178)
25	VSRPGALTRG as for subtype 2f	(SEQ ID NO 179)
	VNQPGALTRG as for subtype 2g	(SEQ ID NO 180)
	VSQPGALTRG as for subtype 2h	(SEQ ID NO 181)
	VSQPGALTKG as for subtype 2i	(SEQ ID NO 182)
	VSRPGALTEG as for subtype 2k	(SEQ ID NO 183)
30	APYIGAPLES or APYTAAPLES as for subtype 4k (S	SEQ ID NO 184 and 185)
	APILSAPLMS as for subtype 41	(SEQ ID NO 186)
	VPNSSVPIHG as for type 9	(SEQ ID NO 187)
	VPNASTPVTG as for subtype 7c	(SEQ ID NO 188)

	VQNASVSIRG as for subtype 7d	(SEQ ID NO 189)
	VKSPCAATAS as for type 10	(SEQ ID NO 190)
	SPRMHHTTQE or SPRLYHTTQE as for sub-	type 1d (SEQ ID NO 191 and 192)
	TSRRHWTVQD as for subtype 1f	(SEQ ID NO 193)
5	APKRHYFVQE as for subtype 2e	(SEQ ID NO 194)
	SPQYHTFVQE as for subtype 2f	(SEQ ID NO 195)
	SPQHHNFSQD as for subtype 2g	(SEQ ID NO 196)
	SPQHHIFVQD as for subtype 2h	(SEQ ID NO 197)
	SPEHHHFVQD as for subtype 2k	(SEQ ID NO 198)
10	RPRRHWTTQD or RPRRHWTAQD or QPRRI	HWTTQD or RPRRHWTTQE as for
	subtype 4k (SEQ ID No	O 199, 200, 201 or 202)
	QPRRHWTVQD as for subtype 41	(SEQ ID NO 203)
	RPKYHQVTQD as for type 9	(SEQ ID NO 204)
	RPRMHQVVQE as for subtype 7c	(SEQ ID NO 205)
15	RPRMYEIAQD as for subtype 7d	(SEQ ID NO 206)
	RHRQHWTVQD as for type 10	(SEQ ID NO 207)
	or a part of said polynucleic acid which is unique to	at least one of the HCV subturns

or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined Table 5, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.

20

25

30

Using the 5' non-coding LiPA system (Stuyver et al., 1993) and a new core LiPA system including multiple probes for subtypes 1a, 1b, 1c, 2a, 2b or 2c derived from the core region (Stuyver et al., 1995), samples from the Benelux, Cameroon, France and Vietnam were selected because of their aberrant reactivities (isolates CAM1078, FR2, FR1, VN4, VN12, VN13, NE98). Some samples were, together with many other samples, sequenced as a control for typing. Sequencing results, however, indicated the discovery of new subtypes (isolates BNL1, BNL2, BNL3, FR4, BNL4, BNL5, BNL6, BNL7, BNL8, BNL9, BNL10, BNL11 and BNL12). Nucleotide sequences in the core and E1 regions which have not yet been reported before, were analyzed in the frame of the invention. Genomic sequences of subtype 1d, 1e, 1f, 1g 2e, 2f, 2g, 2h, 2i, 2k, 2l, 3g, 4k, 4l, 4m, 7a, 7c, 7d and types 9, 10 and 11 isolates are reported for the first time in the present invention. The NS5B region was also analyzed.

The term "polynucleic acid" refers to a single- stranded or double-stranded

nucleic acid sequence which may contain at least 5 contiguous nucleotides in common with the complete nucleotide sequence (e.g. at least 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 60, 75 or more contiguous nucleotides). A polynucleic acid which is up till about 100 nucleotides in length is often also referred to as an oligonucleotide. A polynucleic acid may consist of deoxyribonucleotides or ribonucleotides, nucleotide analogues or modified nucleotides, or may have been adapted for therapeutic purposes. A polynucleic acid may also comprise a double stranded cDNA clone which can be used for cloning purposes, or for *in vivo* therapy, or prophylaxis.

10

5

The oligonucleotides according to the present invention, used as primers or probes may also contain or consist of nucleotide analogous such as phosphorothioates (Matsukura et al., 1987), alkylphosphoriates (Miller et al., 1979) or peptide nucleic acids (Nielsen et al., 1991; Nielsen et al., 1993) or may contain interculating agents (Asseline et al., 1984).

15

As most other variations or modifications introduced into the original DNA sequences of the invention these variations will neccissitate adaptions with respect to the conditions under which the oligonucleotide should be used to obtain the required specificty and sensitivity. However the eventual results will be essentially the same as those obtained with the unmodified oligonucleotides.

20

The introduction of these modifications may be advantageous in order to positivily influence characteristics such as hybridization kinetics, reversibility of the hybrid-formation, biological stability of the oligonucleotide molecules, etc.

The polynucleic acids of the invention may be comprised in a composition of any kind. Said composition may be for diagnostic, therapeutic or prophylactic use.

25

30

The expression "sequences which are unique to an HCV type or subtype" refers to sequences which are not shared by any other type or subtype of HCV, and can thus be used to uniquely detect that HCV type or subtype. Sequence variability is demonstrated in the present invention between the newly found HCV types and subtypes (see Table 5) and the known HCV types and subtypes (see Table 3), and it is therefore from these regions of sequence variability in particular that type- or subtypes-specific polynucleic acids, oligonucleotides, polypeptides and peptides may be obtained. The term type- or subtypes-specific refers to the fact that a sequence is unique to that HCV type or subtype involved.

The expression "nucleotides corresponding to" refers to nucleotides which are homologous or complementary to an indicated nucleotide sequence or region within a specific HCV sequence.

The term "coding region" corresponds to the region of the HCV genome that encodes the HCV polyprotein. In fact, it comprises the complete genome with the exception of the 5' untranslated region and 3' untranslated region.

The term "HCV polyprotein" refers to the HCV polyprotein of the HCV-J isolate (Kato et al., 1990). The adenine residue at position 330 (Kato et al., 1990) is the first residue of the ATG codon that initiates the long HCV polyprotein of 3010 amino acids in HCV-J and other type 1b isolates, and of 3011 amino acids in HCV-1 and other type 1a isolates, and of 3033 amino acids in type 2 isolates HC-J6 and HC-J8 (Okamoto et al., 1992).

This adenine is designated as position 1 at the nucleic acid level, and this methionine is designated as position 1 at the amino acid level, in the present invention. As type 1a isolates contain 1 extra amino acid in the NS5A region, coding sequences of type 1a and 1b have identical numbering in the Core, E1, NS3, and NS4 region, but will differ in the NS5B region as indicated in Table 1. Type 2 isolates have 4 extra amino acids in the E2 region, and 17 or 18 extra amino acids in the NS5 region compared to type 1 isolates, and will differ in numbering from type 1 isolates in the NS3/4 region and NS5b regions as indicated in Table 1. Similar insertions compared with type 1 (but of a different size) can also be observed in type 3a sequences which affect the numbering of type 3a amino acids accordingly. Other insertions or deletions may be readily observed in type1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and 11 sequences after alignment withknown HCV sequences.

25

5

10

15

20

TABLE 1

Region	in	Positions described for HCV-J (Kato et al., 1990)	Positions described for HCV-1 (Choo et al., 1991)	Positions described for HC-J6, HC-J8 (Okamoto et al., 1992)
--------	----	---	---	---

Nucleotides	N\$5B	8023/8235 7932/8271	8352/8564 8261/8600	8026/8238 7935/8274	8433/8645 8342/8681
		coding region of present invention	330/9359	1/9033	342/9439
Amino Acids	NS5B	2675/2745 2645/2757	2675/2745 2645/2757	2676/2746 2646/2758	2698/2768 2668/2780

Table 1:

5

10

Comparison of the HCV nucleotide and amino acid numbering system used in the present invention (*) with the numbering used for other prototype isolates. For example, 8352/8564 indicates the region designated by the numbering from nucleotide 8352 to nucleotide 8564 as described by Kato et al. (1990). Since the numbering system of the present invention starts at the polyprotein initiation site, the 329 nucleotides of the 5' untranslated region described by Kato et al. (1990) have to be substracted, and the corresponding region is numbered from nucleotide 8023 ('8352-329') to 8235 ('8564-329').

The term "genotype" as used in the present invention refers to both types and/or subtypes.

15

20

The term "HCV type" corresponds to a group of HCV isolates of which the complete genome shows more than 73% preferably more than 74% homology at the nucleic acid level, or of which the NS5 region between nucleotide positions 7932 and 8271 shows more than 75.4% homology at the nucleic acid level, or of which the complete HCV polyprotein shows more than 78% homology at the amino acid level, or of which the NS5 region between amino acids at positions 2645 and 2757 shows more than 80% homology at the amino acid level, to polyproteins of the other isolates of the group, with said numbering beginning at the first ATG codon or first methionine of the long HCV polyprotein of the HCV-J isolate (Kato et al., 1990). Isolates belonging to different types of HCV exhibit homologies, over the complete genome, of less than 74%, preferably less than 73%, at the nucleic acid level and less than 78% at the amino acid level. Isolates belonging to the same type usually

25

10

15

20

25

30

show homologies of about 90 to 99% at the nucleic acid level and 95 to 96% at the amino acid level when belonging to the same subtype, and those belonging to the same type but different subtypes preferably show homologies of about 76% to 82% (more particularly of about 77% to 80%) at the nucleic acid level and 85-86% at the amino acid level.

More preferably the definition of HCV types is concluded from the classification of HCV isolates according to their nucleotide distances calculated as detailed below:

(1) based on phylogenetic analysis of nucleic acid sequences in the NS5B region between nucleotides 7935 and 8274 (Choo et al., 1991) or 8261 and 8600 (Kato et al., 1990) or 8342 and 8681 (Okamoto et al., 1991), isolates belonging to the same HCV type show nucleotide distances of less than 0.34, usually less than 0.33, and more usually of less than 0.32, and isolates belonging to the same subtype show nucleotide distances of less than 0.135, usually of less than 0.13, and more usually of less than 0.125, usually ranging between 0.0003 and 0.1151, and consequently isolates belonging to the same type but different subtypes show nucleotide distances ranging from 0.135 to 0.34, usually ranging from 0.1384 to 0.2977, and more usually ranging from 0.15 to 0.32, and isolates belonging to different HCV types show nucleotide distances greater than 0.34, usually greater that 0.35, and more usually of greater than 0.358, more usually ranging from 0.3581 to 0.6670.

(2) based on phylogenetic analysis of nucleic acid sequences in the core/E1 region between nucleotides 378 and 957, isolates belonging to the same HCV type show nucleotide distances of less than 0.38, usually of less than 0.37, and more usually of less than 0.364, and isolates belonging to the same subtype show nucleotide distances of less than 0.17, usually of less than 0.16, and more usually of less than 0.15, more usually less than 0.135, more usually less than 0.134, and consequently isolates belonging to the same type but different subtypes show nucleotide distances ranging from 0.15 to 0.38, usually ranging from 0.16 to 0.37, and more usually ranging from 0.17 to 0.36, more usually ranging from 0.133 to 0.379, and isolates belonging to different HCV types show nucleotide distances greater than 0.34, 0.35, 0.36, usually more than 0.365, and more usually of greater than 0.37,

Table 2: Molecular evolutionary distances

Region	Core/E1	E1	NS5B	NS5B
	579 bp	384 bp	340 bp	222 bp
lsolates*	0.0017 - 0.1347	0.0026 - 0.2031	0.0003 - 0.1151	0.000 - 0.1323
	(0.0750 ± 0.0245)	(0.0969 ± 0.0289)	(0.0637 ± 0.0229)	(0.0607 ± 0.0205)
Subtypes	0.1330 - 0.3794	0.1645 - 0.4869	0.1384 - 0.2977	0.117 - 0.3538
	(0.2786 ± 0.0363)	(0.3761 ± 0.0433)	(0.2219 ± 0.0341)	(0.2391 ± 0.0399)
Types°	0.3479 - 0.6306	0.4309 - 0.9561	0.3581 - 0.6670	0.3457 - 0.7471
	(0.4703 ± 0.0525)	(0.6308 ± 0.0928)	(0.4994 ± 0.0495)	(0.5295 ± 0.0627)

Table 2

Figures created by the PHYLIP program DNADIST are expressed as minimum to maximum (average <u>+</u> standard deviation). Phylogenetic distances for isolates belonging to the same subtype ('isolates'), to different subtypes of the same type ('subtypes'), and to different types ('types') are given.

10

5

In a comparative phylogenetic analysis of available sequences, ranges of molecular evolutionary distances for different regions of the genome were calculated, based on 19,781 pairwise comparisons by means of the DNADIST program of the phylogeny inference package PHYLIP version 3.5c (Felsenstein, 1993). The results are shown in Table 2 and indicate that although the majority of distances obtained in each region fit with classification of a certain isolate, only the ranges obtained in the 340bp NS5B-region are non-overlapping and therefore conclusive. However, as was performed in the present invention, it is preferable to obtain sequence information from at least 2 regions before final classification of a given isolate.

20

15

Designation of a number to the different types of HCV and HCV nomenclature is based on chronological discovery of the different types. The numbering system used in the present invention might still fluctuate according to international conventions or guidelines. For example, "type 4" might be changed into "type 5" or "type 6". Also the arbitrarily chosen border distances between types and subtypes and isolates may still be subject to change according to international guidelines or

conventions. Therefore types 7a, 8a, 8b, 9a may for example be designated 6b, 6c, 6d, and 6d in the future; and type 10a which shows relatedness with genotype 3 may be denoted 3g instead of 10a.

The term "subtype" corresponds to a group of HCV isolates of which the complete polyprotein shows a homology of more than 90% both at the nucleic acid and amino acid levels, or of which the NS5 region between nucleotide positions 7932 and 8271 shows a homology of more than 90% at the nucleic acid level to the corresponding parts of the genomes of the other isolates of the same group, with said numbering beginning with the adenine residue of the initiation codon of the HCV polyprotein. Isolates belonging to the same type but different subtypes of HCV show homologies of more than 74% at the nucleic acid level and of more than 78% at the amino acid level.

It is to be understood that extremely variable regions such as the E1, E2 and NS4 regions will exhibit lower homologies than the average homology of the complete genome of the polyprotein.

Using these criteria, HCV isolates can be classified into at least 11 types. Several subtypes can clearly be distinguished in types 1, 2, 3, 4 and 7: 1a, 1b, 1c, 1d, 1e, 1f, 1g, 2a, 2b, 2c, 2d, 2e, 2f, 2g, 2h, 2i, 2k, 2l, 3a, 3b, 3c, 3d, 3f, 3g, 4a, 4b, 4c, 4d, 4e, 4f, 4g, 4h, 4i, 4j, 4k, 4l, 4m, 7a, 7c, and 7d based on homologies of the 5' UR and coding regions. An overview of most of the reported isolates and their proposed classification according to the typing system of the present invention as well as other proposed classifications is presented in Table 3.

Table 3

5

10

15

20

HCV CLASSIFICATION

25		OKA- MOTO	MORI	СНА	NAKAO	PROTOTYPE
	la	1	1	Pt	GI	HCV-1, HCV-H, HC-JI
	lb	11	II	KI	GII	HCV-J, HCV-BK, HCV-T, HC-JK1, HC-J4, HCV-CHINA
	lc					HC-G9
	2a	111	III	K2a	GIII	HC-J6
30	2b	rv	īv	K2b	GIII	HC-J8

	2c					S83, ARG6, ARG8, I10, T983
	2d					NE92
	3a	v	v	К3	GIV	BR36, BR56, HD10, N2L1, BR33, Ta, E-b1
5	3Ь		VI	K3	GIV	HCV-TR, Tb, NE137
	3c					NE48
	3d					NE274
	3e					NE145
	3f					NE125
10	4a					Z4, GB809-4
	4b					Z 1
	4c					GB116, GB358, GB215, Z6, Z7
	4d					DK13
	4c					GB809-2, CAM600, CAM736
15	4f					CAM622, CAM627
	4g					GB549
	4h					GB438
	4i					CAR4/1205
	4j					CAR1/905
20	5a				GV	SA3, SA4, SA1, SA7, SA11, BE95
	6a					HK1, HK2, HK3, HK4, VN11

<u>Table 3</u> Overview of the known HCV types and subtypes classified according to the different authors.

The term "complement" refers to a nucleotide sequence which is complementary to an indicated sequence and which is able to hybridize to the indicated sequences.

The composition of the invention can comprise many combinations. By way of example, the composition of the invention can comprise:

- two (or more) nucleic acids from the same region or,
- two nucleic acids (or more), respectively from different regions, for the same isolate or for different isolates,
 - or nucleic acids from the same regions and from at least two different regions

(for the same isolate or for different isolates).

5

10

15

20

25

30

The present invention relates particularly to a polynucleic acid as defined above having a sequence selected from any of SEQ ID NO 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103 to 105, or a part of said polynucleic acid which is unique to any of the HCV subtypes or types as defined in Table 5, and which contains at least one nucleotide differing from known HCV polynucleic acids, or the complement thereof.

The present invention relates more particularly to a polynucleic acid as defined above, which codes for the 5' UR, the Core/E1, the NS4 or the NS5B region or a part thereof.

More particularly, the present invention relates to a polynucleic acid as defined above which is a cDNA sequence.

Also included within the present invention are sequence variants of the polynucleic acids as selected from any of the nucleotide sequences as given in any of the above given SEQ ID numbers with said sequence variants containing either deletion and/or insertions of one or more nucleotides, especially insertions or deletions of 1 or more codons, mainly at the extremities of oligonucleotides (either 3' or 5'), or substitutions of some non-essential nucleotides (i.e. nucleotides not essential to discriminate between different genotypes of HCV) by others (including modified nucleotides an/or inosine), for example, a type 1 or 2 sequence might be modified into a type 7 sequence by replacing some nucleotides of the type 1 or 2 sequence with type-specific nucleotides of type 7 as shown in for instance Figure 1 and 2.

Particularly preferred variant polynucleic acids of the present invention include also sequences which hybridise under stringent conditions with any of the polynucleic acid sequences of the present invention. Particularly, sequences which show a high degree of homology (similarity) to any of the polynucleic acids of the invention as described above. Particularly sequences which are at least 80%, 85%, 90%, 95% or more homologous to said polynucleic acid sequences of the invention. Preferably said sequences will have less than 20%, 15%, 10%, or 5% variation of the original nucleotides of said polynucleic acid sequence.

Polynucleic acid sequences according to the present invention which are

10

15

20

25

30

homologous to the sequences as represented by a SEQ ID NO can be characterized and isolated according to any of the techniques known in the art, such as amplification by means of sequence-specific primers, hybridization with sequence-specific probes under more or less stringent conditions, serological screening methods or via the LiPA typing system.

Other preferred variant polynucleic acids of the present invention include sequences which are redundant as a result of the degeneracy of the genetic code compared any of the above-given polynucleic acids of the present invention. These variant polynucleic acid sequences will thus encode the same amino acid sequence as the polynucleic acids they are derived from.

Also included within the scope of the present invention are 5' non-coding region sequences which can be readily obtained from type 1 subtype 1d, 1e, 1f or 1g isolates; type 2 subtype 2e, 2f, 2g, 2h, 2i, 2k or 2l isolates; type 3 subtype 3g isolates; type 4 subtype 4k, 4l or 4m isolates; type 7 subtype 7a, 7c or 7d isolates, type 9, type 10 or type 11 isolates discribed herein. Such sequences may contain type or subtype-specific motifs which can be employed for type and/or subtype-specific hybridization assays, e.g. such as described by Stuyver et al. (1993).

Polynucleic acid sequences of the genomes indicated above from regions not yet depicted in the present examples, figures and sequence listing can be obtained by any of the techniques known in the art, such as amplification techniques using suitable primers from the sequences of these new genomes given in Figure 1 of the present invention.

The present invention also relates to an oligonucleotide primer comprising part of a polynucleic acid as defined above, with said primer being able to act as a primer for specifically amplifying the nucleic acid of a certian HCV isolate belonging to the genotype from which the primer is derived.

The term "primer" refers to a single stranded DNA oligonucleotide sequence capable of acting as a point of initiation for synthesis of a primer extension product which is complementary to the nucleic acid strand to be copied. The length and the sequence of the primer must be such that they allow to prime the synthesis of the extension products. Preferably the primer is about 5-50 nucleotides. Specific length and sequence will depend on the complexity of the required DNA or RNA targets, as well as on the conditions of primer use such as temperature and ionic strength.

10

15

20

25

30

The fact that amplification primers do not have to match exactly with corresponding template sequence to warrant proper amplification is amply documented in the literature (Kwok et al., 1990).

The amplification method used can be either polymerase chain reaction (PCR; Saiki et al., 1988), ligase chain reaction (LCR; Landgren et al., 1988; Wu & Wallace, 1989; Barany, 1991), nucleic acid sequence-based amplification (NASBA; Guatelli et al., 1990; Compton, 1991), transcription-based amplification system (TAS; Kwoh et al., 1989), strand displacement amplification (SDA; Duck, 1990; Walker et al., 1992) or amplification by means of Qß replicase (Lizardi et al., 1988; Lomeli et al., 1989) or any other suitable method to amplify nucleic acid molecules using primer extension. During amplification, the amplified products can be conveniently labelled either using labelled primers or by incorporating labelled nucleotides. Labels may be isotopic (32P, 35S, etc.) or non-isotopic (biotin, digoxigenin, etc.). The amplification reaction is repeated between 20 and 70 times, advantageously between 25 and 45 times.

The present invention also relates to an oligonucleotide probe comprising part of a polynucleic acid as defined above, with said probe being able to act as a hybridization probe for specific detection and/or classification into types and/or subtypes of an HCV nucleic caid containing said nucleotide sequence, with said probe being possibly labelled or attached to a solid substrate.

The term "probe" refers to single stranded sequence-specific oligonucleotides which have a sequence which is complementary to the target sequence of the HCV genotype(s) to be detected.

Preferably, these probes are about 5 to 50 nucleotides long, more preferably from about 10 to 25 nucleotides.

The term "solid support" can refer to any substrate to which an oligonucleotide probe can be coupled, provided that it retains its hybridization characteristics and provided that the background level of hybridization remains low. Usually the solid substrate will be a microtiter plate, a membrane (e.g. nylon or nitrocellulose) or a microsphere (bead). Prior to application to the membrane or fixation it may be convenient to modify the nucleic acid probe in order to facilitate fixation or improve the hybridization efficiency. Such modifications may encompass homopolymer tailing, coupling with different reactive groups such as aliphatic

10

15

25

groups, $\mathrm{NH_2}$ groups, SH groups, carboxylic groups, or coupling with biotin or haptens.

The present invention also relates to a diagnostic kit for use in determining the genotype of HCV, said kit comprising a primer as defined above.

The present invention also relates to a diagnostic kit for use in determining the genotype of HCV, said kit comprising a probe as defined above.

The present invention also relates to a diagnostic kit as defined above, wherein said probe(s) is(are) attached to a solid substrate.

The present invention also relates to a diagnostic kit as defined above, wherein a range of said probes is attached to specific locations on a solid substrate.

The present invention also relates to a diagnostic kit as defined above, wherein said solid support is a membrane strip and said probes are coupled to the membrane in the form of parallel lines.

The present invention also relates to a method for the detection of HCV nucleic acids present in a biological sample, comprising:

- (i) possibly extracting sample nucleic acid,
- (ii) amplifying the nucleic acid with at least one primer as defined above,
- (iii) detecting the amplified nucleic acids.

The present invention also relates to a method for the detection of HCV nucleic acids present in a biological sample, comprising:

- (i) possibly extracting sample nucleic acid,
- (ii) possibly amplifying the nucleic acid with at least one primer as defiend above, or with a universal HCV primer,
- (iii) hybridizing the nucleic acids of the biological sample, possibly under denatured conditions, at appropriate conditions with one or more probes as defined above, with said probes being preferably attached to a solid substrate,
- (iv) possibly washing at appropriate conditions,
- (v) detecting the hybrids formed.

The present invention also relates to a method for detecting the presence of one or more HCV genotypes present in a biological sample, comprising:

- (i) possibly extracting sample nucleic acid,
- (ii) specifically amplifying the nucleic acid with at least one primer as defined

15

20

25

30

above.

(iii) detecting said amplified nucleic acids.

The present invention also relates to a method for detecting the presence of one or more HCV genotypes present in a biological sample, comprising:

- 5 (i) possibly extracting sample nucleic acid,
 - (ii) possibly amplifying the nucleic acid with at least one primer as defined above or with a universal HCV primer,
 - hybridizing the nucleic acids of the biological sample, possibly under denatured conditions, at appropriate conditions with one or more probes as defined above, with said probes being preferably attached to a solid substrate,
 - (iv) possibly washing at appropriate conditions,
 - (v) detecting the hybrids formed.
 - (vi) inferring the presence of one or more HCV genotypes present from the observed hybridization pattern.

The present invention also relates to a method as defined above, wherein said probes are further characterized as defined above.

The present invention also relates to a method as defined above, wherein said nucleic acids are labelled during or after amplification.

Preferably, this technique could be performed in the 5' non-coding, Core or NS5B region.

The term "nucleic acid" can also be referred to as analyte strand and corresponds to a single- or double-stranded nucleic acid molecule. This analyte strand is preferentially positive- or negative stranded RNA, cDNA or amplified cDNA.

The term "biological sample" refers to any biological sample (tissue or fluid) containing HCV nucleic acid sequences and refers more particularly to blood serum or plasma samples.

The term "universal HCV primer" refers to oligonucleotide sequences complementary to any of the conserved regions of the HCV genome.

The expression "appropriate" hybridization and washing conditions are to be understood as stringent and are generally known in the art (e.g. Maniatis et al., Molecular Cloning: A Laboratory Manual, New York, Cold Spring Harbor Laboratory, 1982).

10

15

20

25

30

However, according to the hybridization solution (SSC, SSPE, etc.), these probes should be hybridized at their appropriate temperature in order to attain sufficient specificity.

The term "labelled" refers to the use of labelled nucleic acids. This may include the use of labelled nucleotides incorporated during the polymerase step of the amplification such as illustrated by Saiki et al. (1988) or Bej et al. (1990) or labelled primers, or by any other method known to the person skilled in the art.

The process of the invention comprises the steps of contacting any of the probes as defined above, with one of the following elements:

 either a biological sample in which the nucleic acids are made available for hybridization,

- or the purified nucleic acids contained in the biological sample
- or a single copy derived from the purified nucleic acids,
- or an amplified copy derived from the purified nucleic acids, with said elements or with said probes being attached to a solid substrate.

The expression "inferring the presence of one or more HCV genotypes present from the observed hybridization pattern" refers to the identification of the presence of HCV genomes in the sample by analyzing the pattern of binding of a panel of oligonucleotide probes. Single probes may provide useful information concerning the presence or absence of HCV genomes in a sample. On the other hand, the variation of the HCV genomes is dispersed in nature, so rarely is any one probe able to identify uniquely a specific HCV genome. Rather, the identity of an HCV genotype may be inferred from the pattern of binding of a panel of oligonucleotide probes, which are specific for (different) segments of the different HCV genomes. Depending on the choice of these oligonucleotide probes, each known HCV genotype will correspond to a specific hybridization pattern upon use of a specific combination of probes. Each HCV genotype will also be able to be discriminated from any other HCV genotype amplified with the same primers depending on the choice of the oligonucleotide probes. Comparison of the generated pattern of positively hybridizing probes for a sample containing one or more unkown HCV sequences to a scheme of expected hybridization patterns, allows one to clearly infer the HCV genotypes present in said sample.

The present invention thus relates to a method as defined above, wherein one SUBSTITUTE SHEET (RULE 26)

or more hybridization probes are selected from any of SEQ ID NO 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103 or 105 or sequence variants thereof as defined above.

5

In order to distinguish the amplified HCV genomes from each other, the target polynucleic acids are hybridized to a set of sequence-specific DNA probes targetting HCV genotypic regions (unique regions) located in the HCV polynucleic acids.

Most of these probes target the most type- or subtype-specific regions of HCV genotypes, but some can be caused to hybridize to more than one HCV genotype.

10

According to the hybridization solution (SSC, SSPE, etc.), these probes should be stringently hybridized at their appropriate temperature in order to attain sufficient specificity. However, by slightly modifying the DNA probes, either by adding or deleting one or a few nucleotides at their extremities (either 3' or 5'), or substituting some non-essential nucleotides (i.e. nucleotides not essential to discriminate between types) by others (including modified nucleotides or inosine) these probes or variants thereof can be caused to hybridize specifically at the same hybridization conditions (i.e. the same temperature and the same hybridization solution). Also changing the amount (concentration) of probe used may be beneficial to obtain more specific hybridization results. It should be noted in this context, that probes of the same length, regardless of their GC content, will hybridize specifically at approximately the same temperature in TMACI solutions (Jacobs et al., 1988).

20

15

Suitable assay methods for purposes of the present invention to detect hybrids formed between the oligonucleotide probes and the nucleic acid sequences in a sample may comprise any of the assay formats known in the art, such as the conventional dot-blot format, sandwich hybridization or reverse hybridization. For example, the detection can be accomplished using a dot blot format, the unlabelled amplified sample being bound to a membrane, the membrane being incorporated with at least one labelled probe under suitable hybridization and wash conditions, and the presence of bound probe being monitored.

30

25

An alternative and preferred method is a "reverse" dot-blot format, in which the amplified sequence contains a label. In this format, the unlabelled oligonucleotide probes are bound to a solid support and exposed to the labelled sample under appropriate stringent hybridization and subsequent washing conditions. It is to be

10

15

20

25

30

understood that also any other assay method which relies on the formation of a hybrid between the nucleic acids of the sample and the oligonucleotide probes according to the present invention may be used.

According to an advantageous embodiment, the process of detecting one or more HCV genotypes contained in a biological sample comprises the steps of contacting amplified HCV nucleic acid copies derived from the biological sample, with oligonucleotide probes which have been immobilized as parallel lines on a solid support.

According to this advantageous method, the probes are immobilized in a Line Probe Assay (LiPA) format. This is a reverse hybridization format (Saiki et al., 1989) using membrane strips onto which several oligonucleotide probes (including negative or positive control oligonucleotides) can be conveniently applied as parallel lines.

The invention thus also relates to a solid support, preferably a membrane strip, carrying on its surface, one or more probes as defined above, coupled to the support in the form of parallel lines.

The LiPA is a very rapid and user-friendly hybridization test. Results can be read after 4 hours, after the start of the amplification. After amplification during which usually a non-isotopic label is incorporated in the amplified product, and alkaline denaturation, the amplified product is contacted with the probes on the membrane and the hybridization is carried out for about 1 to 1,5 h hybridized polynucleic acid is detected. From the hybridization pattern generated, the HCV type can be deduced either visually, but preferably using dedicated software. The LiPA format is completely compatible with commercially available scanning devices, thus rendering automatic interpretation of the results very reliable. All those advantages make the LiPA format liable for the use of HCV detection in a routine setting. The LiPA format should be particularly advantageous for detecting the presence of different HCV genotypes.

The present invention also relates to a method for detecting and identifying novel HCV genotypes, different from the known HCV genomes, comprising the steps of:

- determining to which HCV genotype the nucleotides present in a biological sample belong, according to the process as defined above,
- in the case of observing a sample which does not generate a hybridization

pattern compatible with those defined in Table 3, sequencing the portion of the HCV genome sequence corresponding to the aberrantly hybridizing probe of the new HCV genotype to be determined.

The present invention also relates to a method for preparing a polynucleic acid according to the present invention. These methods include any method known in the art for preparing polynucleic acids (e.g. the phosphodiester method for synthesizing oligonucleotides as described by Agarwal et al. 1972, Agnew. Chem. Int. Ed. Engl. 11:451, the phosphotriester method of Hsiung et al. 1979, Nucleic Acid Res. 6:1371, or the automated diethylphosphoramidite method of Baeucage et al. 1981, Tetrahedron Letters 22:1859-1862.). Alternatively, the polynucleic acids of the present invention may be isolated fragments of naturally occuring or cloned DNA or RNA. In addition, the oligonucleotides according to the present invention may be synthesized automatically on commercial instruments sold by a variety of manufacturers.

15

10

5

The present invention particularly also relates to a polypeptide having an amino acid sequence encoded by a polynucleic acid as defined above, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in Table 5, and which contains at least one amino acid differing from any of the known HCV types or subtypes, or an analog thereof being substantially homologous and biologically equivalent.

20

25

The term 'polypeptide' refers to a polymer of amino acids and does not refer to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. Included within the definition are, for example, polypeptides containing one or more analogues of an amino acid (including, for example, unnatural amino acids, PNA, etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

30

The term "unique" is referred above.

By "biologically equivalent" as used throughout the specification and claims, it is meant that the compositions are immunogenically equivalent to the proteins (polypeptides) or peptides of the invention as defined above and below.

10

15

20

25

30

By "substantially homologous" as used throughout the ensuing specification and claims to describe proteins and peptides, it is meant a degree of homology in the amino acid sequence to the proteins or peptides of the invention. Preferably the degree of homology is in excess of 90, preferably in excess of 95, with a particularly preferred group of proteins being in excess of 99 homologous with the proteins or peptides of the invention.

The term "analog" as used throughout the specification or claims to describe the proteins or peptides of the present invention, includes any protein or peptide having an amino acid residue sequence substantially identical to a sequence specifically shown herein in which one or more residues have been conservatively substituted with a biologically equivalent residue. Examples of conservative substitutions include the substitution of one-polar (hydrophobic) residue such as isoleucine, valine, leucine or methionine for another, the substitution of one polar (hydrophillic) residue for another such as between arginine and lysine, between glutamine and asparagine, between glycine and serine, the substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another. Examples of allowable mutations according to the present inevntion can be found in Table 4.

The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that the resulting protein or peptide is biologically equivalent to the protein or peptide of the invention.

"Chemical derivative" refers to a protein or peptide having one or more residues chemically derivatized by reaction of a functional side group. Examples of such derivatized molecules, include but are not limited to, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloracetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-imbenzylhistidine. Also included as chemical derivatives are those proteins or peptides which contain one or more naturally-occurring amino acid derivatives of the twenty standard amino acids. For examples: 4-hydroxyproline may be substituted for lysine; 3-

10

15

20

25

methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine. The proteins or peptides of the present invention also include any protein or peptide having one or more additions and/or deletions or residues relative to the sequence of a peptide whose sequence is shown herein, so long as the peptide is biologically equivalent to the proteins or peptides of the invention.

It is to be noted that, at the level of the amino acid sequence, at least one amino acids difference (with respect to known HCV amino acid sequences) is sufficient to be part of the invention, which means that the polypeptides of the invention correspond to polynucleic acids having at least one nucleotide difference (with known HCV polynucleic acid sequences) involving an amino acid difference in the encoded polyprotein.

As the NS4 and the Core regions are known to contain several epitopes, for example characterized in patent application EP-A-O 489 968, and as the E1 protein is expected to be subject to immune attack as part of the viral envelope and expected to contain epitopes, the NS4, Core and E1 epitopes of the new types and subtypes disclosed herein will consistently differ from the epitopes present in previously known genotypes. This is examplified by the type-specificity of NS4 synthetic peptides as described in Simmonds et al. (1993c) and Stuyver et al. (1993b) and PCT/EP 94/O1323 and the type-specificity of recombinant E1 proteins as described in Maertens et al. (1994).

The peptides according to the present invention contain preferably at least 3, preferably 4, 5 contiguous HCV amino acids, 6, 7 preferably however at least 8 contiguous HCV amino acids, at least 10 or at least 15 (for instance at least 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50 or more amino acids).

TABLE 4

30	Amino acids	Synonymous groups
	Ser (S)	Ser, Thr, Gly, Asn
	Arg (R)	Arg, His, Lys, Glu, Gln

25

30

35

		20		
5	Leu (L) Pro (P) Thr (T) Ala (A) Val (V) Gly (G) Ile (I) Phe (F) Tyr (Y)	Leu; Ile, Met, Phe, Val, Tyr Pro, Ala, Thr, Gly Thr, Pro, Ser, Ala, Gly, His, Gln Ala, Pro, Gly, Thr Val, Met, Ile, Tyr, Phe, Leu, Val Gly, Ala, Thr, Pro, Ser Ile, Met, Leu, Phe, Val, Ile, Tyr Phe, Met, Tyr, Ile, Leu, Trp, Val Tyr, Phe, Trp, Met, Ile, Val, Leu Cys, Ser, Thr, Met		
15	Cys (C) His (H) Gln (Q) Asn (N) Lys (K) Asp (D) Glu (E) Met (M)	His, Gln, Arg, Lys, Glu, Thr Gln, Glu, His, Lys, Asn, Thr, Arg Asn, Asp, Ser, Gln Lys, Arg, Glu, Gln, His Asp, Asn, Glu, Gln Glu, Gln, Asp, Lys, Asn, His, Arg Met, Ile, Leu, Phe, Val		

<u>Table 4</u> Overview of the amino acid substitutions which could form the basis of analogs (muteins) as defined above

The polypeptides of the invention, and particularly the fragments, can be prepared by classical chemical synthesis.

The synthesis can be carried out in homogeneous solution or in solid phase.

For instance, the synthesis technique in homogeneous solution which can be used is the one described by Houbenweyl in the book entitled "Methode der organischen chemie" (Method of organic chemistry) edited by E. Wunsh, vol. 15-I et II. THIEME, Stuttgart 1974.

The polypeptides of the invention can also be prepared in solid phase according to the methods described by Atherton and Shepard in their book entitled "Solid phase peptide synthesis" (IRL Press, Oxford, 1989).

The polypeptides according to this invention can be prepared by means of recombinant DNA techniques as described by Maniatis et al., Molecular Cloning: A Laboratory Manual, New York, Cold Spring Harbor Laboratory, 1982).

The present invention relates particularly to a polypeptide as defined above, comprising in its amino acid sequence at least one of the following amino acid residues:

10

15

20

25

30

115, C38, V44, A49, Q43, P49, Q55, A58, S60 or D60, E68 or V68, H70, A71 or Q71 or N71, D72, H81, H101, D106, S110, L130, I134, E135, L140, S148, T150 or E150, Q153, F155, D157, G160, E165, I169, F181, L186, T190, T192 or I192 or H192, I193, A195, S196, R197 or N197 or K197, Q199 or D199 or H199, N199, F200 or T200, A208, I213, M216 or S216, N217 or S217 or G217 or K217, T218, I219, A222, Y223, I230, W231 or L231, S232 or H232 or A232, Q233, E235 or L235, F236 or T236, F237, L240 or M240, A242, N244, N249, I250 or K250 or R250, A252 or C252, A254, I255 or V255, D256 or M256, E257, E260 or K260, R261, V268, S272 or R272, I285, G290 or F290, A291, A293 or L293 or W293, T294 or A294, S295, H295, K296 or E296, Y297 or M297, I299 or Y299, I300, S301, P316, S2646, A2648, G2649, A2650, V2652, Q2653, H2656 or L2656, D2657, F2659, K2663 or Q2663, A2667 or V2667, D2677, L2681, M2686 or Q2686 or E2686, A2692 or K2692, H2697, I2707, L2708 or Y2708, A2709, A2719 or M2719, F2727, T2728 or D2728, E2729, F2730 or Y2730, 12741, 12745, V2746 or E2746 or L2746 or K2746, A2748, S2749 or P2749, R2750, E2751, D2752 or N2752 or S2752 or T2752 or V2752 or I2752 or Q2752, S2753 or D2753 or G2753, D2754, A2755, L2756 or Q2756, or R2757,

with said notation being composed of a letter representing the amino acid residue by its one-letter code, and a number representing the amino acid numbering according to Kato et al., 1990 as shown in Table 1 (see also the numbering in Figures 2, 4 and 6),

or a part thereof which is unique to at least one of the HCV subtypes or types as defined in Table 5, and which contains at least one amino acid differing from any of the known HCV types or subtypes, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide or part thereof.

These unique amino acid residues can be deduced from aligning the new HCV amino acid sequences as given in Figure 3 to all known HCV sequences. An alignment with the new sequences as represented in SEQ ID NO 1 to 106 is given in for instance Figures 2, 4 and 6. It should be clear that the alignments given in these figures may be completed with all known HCV sequences to illustrate that any of the above-given unique residues is indeed unique for at least one of the new HCV sequences of the present invention.

Within the group of unique and new amino acid residues of the present

10

15

20

25

30

invention, unique residues may be found which are specific for the following new types (subtypes) of HCV according to the HCV classification system used in the present invention: type 1 subtype 1d, 1e, 1f or 1g isolates; type 2 subtype 2e, 2f, 2g, 2h, 2i, 2k or 2l isolates; type 3 subtype 3g isolates; type 4 subtype 4k, 4l or 4m isolates; type 7 subtype 7a, 7c or 7d isolates, type 9, type 10 or type 11 isolates. In order to obtain these residues the alignments given in Figures 2, 4 and 6 may be used to deduce the type- and or subtype-specificity of any of the unique residues given above.

For example T190 (detected in subtype 1d) refers to a threomine at position 190 (see Figure 2). In other sequences only a serine (S190) or exceptionally an alanine (A190 in type 10a) can be detected.

The polypeptides according to this embodiment of the invention may be possibly labelled, or attached to a solid substrate, or coupled to a carrier molecule such as biotin, or mixed with a proper adjuvant all known in the art and according to the intended use (diagnostic, therapeutic or prophylactic).

The present invention also relates to a polypeptide as defined above, comprising in its amino acid sequence at least one of the sequences repesented by SEQ ID NO107 to 207 as listed above, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in Table 5, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide or part thereof.

The present invention relates also to a polypeptide having an amino acid sequence as represented in any of SEQ ID NO 1 to 106, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in Table 5, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide or part thereof.

The variable region in the core protein (V-CORE in Fig. 2) has been shown to be useful for serotyping (Machida et al., 1992). The sequence of the type 1 subtype 1d, 1e, 1f or 1g sequence; type 2 subtype 2e, 2f, 2g, 2h, 2i, 2k and 2l sequence; type 3 subtype 3g; type 4, subtype 4k, 4l or 4m sequence; type 7 (subtype 7a, 7c and 7d sequences), 9, 10 or 11 sequences of the present invention show type-specific features in this region. The peptide from amino acid 68 to 78 (V-core region) shows the following unique sequence for the sequences of the present invention (see

	figure 2):		
	ARQSDGRSWAQ or ARRSEGRSWAQ as for subtype 1d (SEQ ID NO 107 and		
	108)		
	ERRPEGRSWAQ as for subtype 1e	(SEQ ID NO	O 109)
5	ARRPEGRSWAQ as for subtype 1f	(SEQ ID NO	O 110)
	DRRTTGKSWGR as for subtype 2k	(SEQ ID NO	0 111)
	DRRATGRSWGR as for subtype 2e	(SEQ ID NO) 112)
	DRRATGKSWGR as for subtype 2f	(SEQ ID NO) 113)
	VRQPTGRSWGQ as for type 9	(SEQ ID NO	114)
10	VRHQTGRTWAQ as for subtype 7a and 7c	(SEQ ID NO	115)
	VRQNQGRTWAQ as for subtype 7d	(SEQ ID NO	116)
	ARRTEGRSWAQ as for type 10	(SEQ	ID NO 117)
	VRRTTGRXXXX or VRRTTGRTWAQ as for to	ype 11	(SEQ ID NO 118 and
	119)		
15	Five type-specific variable regions (V1 to V5) can be ider	ntified after aligning
	E1 amino acid sequences of the genotypes of the pre	sent inventio	on to the genotypes
	already known, as shown in Figure 2.		
	Region V1 encompasses amino acids 192 to	203, this is	the amino-terminal
	10 amino acids of the E1 p	protein. The	following unique
20	sequences as shown in Fig. 2 can be deduced:		
	HEVRNASGVYHV or HEVRNASGVYHL as for	subtype 1d,	(SEQ ID NO
	120 and 121)		
	YEVHSTTDGYHV as for subtype 1f	(SEQ ID NO	122)
		(SEQ ID NO	123)
25		(SEQ ID NO	124)
		SEQ ID NO	125)
		SEQ ID NO	126)
		SEQ ID NO	•
	VEIKNTXNTYVL or VEIKNTSNTYVL as for sub	otype 2k	(SEQ ID NO 128
30	and 129)		
INYRNVSGIYYV or INYRNTSGIYHV or INYHNTSGIYHI or TNYRNVSGIYH			NYRNVSGIYHV a
	for subtype 4k (SEQ ID NO 130, 131, 132 or 133)		
	QHYRNVSGIYHV as for subtype 4I (SEQ ID	NO 134)	

	30			
	IQVKNASGIYHL as for type 9 (SEQ ID NO 135)			
	AHYTNKSGLYHL as for subtype 7c (SEQ ID NO 136)			
	LNYANKSGLYHL as for subtype 7d (SEQ ID NO 137)			
	(SEQ ID NO 138)			
5	Region V2 encompasses amino acids 213 to 223. The following unique			
J	requences can be found in the V2 region as shown in Figure 2:			
IYEMDGMIMHY or IYEMSGMILHA as for subtype 1d, (SEQ ID NO 139				
	and 140) (SEQ ID NO 141)			
10	VAEVKDIITH! as lot amprahe !!			
	VWQLXDAVLHV as for subtype 2e (SEQ ID NO 142)			
	VWQLRDAVLHV as for subtype 2f (SEQ ID NO 143)			
	IWQMQGAVLHV as for subtupe 2g (SEQ ID NO 144)			
	VWQLKDAVLHV as for subtype 2h (SEQ ID NO 145)			
	VWQLEEAVLHV as for subtype 2i (SEQ ID NO 146)			
15	TWQLXXAVLHV as for subtype 2k (SEQ ID NO 147)			
*	VYEADHHILHL or VYEADHHILAL or VFEADHHILHL as for subtype 4k			
(SEQ ID NO 148, 149 and 150) (SEQ ID NO 151)				
	VYESDHHILHL as for subtype 4			
	VEENE IMILIAL as for type o			
20	VYEAETLILHL as for subtype 70			
	VYEANGMILML as for subtype 7.5			
Region V3 encompasses the amino acids 230 to 242. The following ur				
V3 region sequences can be deduced from Figure 2:				
25	VREDNHLRCWMAL or VRENNSSRCWMAL as for subtype 1d			
	(SEQ ID NO 156 and 157) (SEQ ID NO 158)			
	IREGNISHOW VLF as 101 30057 FF			
	ENSSGRENCEWIFT as 10. sectors			
	ERSGNRIFCWIAV as 10. 5555/F5			
30	ELQGNKSRCWIFV as for 3333775 -3			
	ERHQNQSRCWIFV as 101 525 775			
	EWKDN1SRCWIFV as to substype and the substype and the substype as the substype			
	EREGNSSRCWIPV as for subtype 2k (SEQ ID NO 165)			

	•) I		
	VREGNQSRCWVAL or VRTGNO	DSRCWVAL or VRVGNQSSCWVAL or		
	VRVGNQSRCWVAL or VKEGNHS			
	(SEQ ID NO 166, 16	7, 168 or 169)		
	VKTGNTSRCWVAL as for subtype 4I (SEQ ID NO 170)			
5	IKAGNESRCWLPV as for type 9	(SEQ ID NO 171)		
	VKXXNQSRCWVQA as for subtyp	e 7c (SEQ ID NO 172)		
	VKTGNLTKCWLSA as for subtype 7d (SEQ ID NO 173)			
	VRSGNTSRCWIPV as for type 10	(SEQ ID NO 174)		
	Region V4 encompasses the amino	acids 248 to 257. The following unique		
10 V4 region sequences can be deduce				
	VKNASVPTAA or VKDANVPTAA as for s	ubtype 1d (SEQ ID NO 175 and 176)		
	ARIANAPIDE as for subtype 1f			
	VSKPGALTKG as for subtype 2e	(SEQ ID NO 178)		
	VSRPGALTRG as for subtype 2f			
15	VNQPGALTRG as for subtype 2g	(SEQ ID NO 180)		
	VSQPGALTRG as for subtype 2h	(SEQ ID NO 181)		
	VSQPGALTKG as for subtype 2i			
	VSRPGALTEG as for subtype 2k			
	APYIGAPLES or APYTAAPLES as fo			
20	and 185)			
	APILSAPLMS as for subtype 4I	(SEQ ID NO 186)		
	VPNSSVPIHG as for type 9	(SEQ ID NO 187)		
	VPNASTPVTG as for subtype 7c	(SEQ ID NO 188)		
	VQNASVSIRG as for subtype 7d	(SEQ ID NO 189)		
25	VKSPCAATAS as for type 10	(SEQ ID NO 190)		
	Region V5 encompasses the amino a	acids 294 to 303. The following unique		
	V5 region peptides can be deduced from fig			
	SPRMHHTTQE or SPRLYHTTQE as for	or subtype 1d (SEQ ID NO 191		
	and 192)			
30	TSRRHWTVQD as for subtype 1f	(SEQ ID NO 193)		
	APKRHYFVQE as for subtype 2e	(SEQ ID NO 194)		
	SPQYHTFVQE as for subtype 2f	(SEQ ID NO 195)		
	SPQHHNFSQD as for subtype 2g	(SEQ ID NO 196)		
	_	- · - · ,		

15

20

25

30

(SEQ ID NO 197) SPQHHIFVQD as for subtype 2h (SEQ ID NO 198) SPEHHHFVQD as for subtype 2k RPRRHWTTQD or RPRRHWTAQD or QPRRHWTTQD or RPRRHWTTQE as for (SEQ ID NO 199, 200, 201 or 202) subtype 4k (SEQ ID NO 203) QPRRHWTVQD as for subtype 41 5 (SEQ ID NO 204) RPKYHQVTQD as for type 9 (SEQ ID NO 205) RPRMHQVVQE as for subtype 7c (SEQ ID NO 206) RPRMYEIAQD as for subtype 7d (SEQ ID NO 207) RHRQHWTVQD as for type 10

The above given list of peptides are particularly useful for treatment and vaccine and diagnostic development.

Also comprised in the present invention is any synthetic peptide (see below) or polypeptide containing at least an epitope derived from the above-defined peptides in their peptidic chain. Also comprised within the present invention is any synthetic peptide or polypeptide comprising at least 6, 7, 8, or 9 contiguous amino acids derived from the above-defined peptides in their peptidic chain.

As used herein, 'epitope' or 'antigenic determinant' means an amino acid sequence that is immunoreactive. Generally an epitope consists of at least 3 to 4 amino acids, and more usually, consists of at least 5 or 6 amino acids, sometimes the epitope consists of about 7 to 8, or even about 10 amino acids.

The present invention particularly relates to any peptide (see below) or polypeptide contained in any of the amino acid sequences as represented in SEQ ID NO 2, 4, 7, 9, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104 or 106 (see Table 5 and Figure 3, Examples section).

The present invention also relates to a recombinant polypeptide encoded by a polynucleic acid as defined above, or a part thereof which is unique to any of the HCV subtypes or types as defined in Table 5, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.

The present invention also relates to a recombinant expression vector comprising a polynucleic acid or a part thereof as defined above, operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements.

10.

15

20

25

30

In general said recombinant vector will comprise a vector sequence, an appropriate prokaryotic, eukaryotic or viral promoter sequence followed by the nucleotide sequences as defined above, with said recombinant vector allowing the expression of any one of the polypeptides as defined above in a prokaryotic, or eukaryotic host or in living mammals when injected as naked DNA, and more particularly a recombinant vector allowing the expression of any of the new HCV sequences of the invention spanning particularly the following amino acid positions:

- a polypeptide starting in the region between positions 1 and 10 and ending at any position in the region between positions 70 and 420, more particularly a polypeptide spanning positions 1 to 70, 1 to 85, positions 1 to 120, positions 1 to 150, positions 1 to 191, or positions 1 to 200, for expression of the Core protein, and a polypeptide spanning positions 1 to 263, positions 1 to 326, positions 1 to 383, or positions 1 to 420 for expression of the Core and E1 protein;
- a polypeptide starting at any position in the region between positions 117 and 192, and ending at any position in the region between positions 263 and 420, for expression of E1, or forms that have the hydrophobic region deleted (positions 264 to 293 plus or minus 8 amino acids);
- a polypeptide starting at any position in the region between positions 1556 and 1688, and ending at any position in the region between positions 1739 and 1764, for expression of NS4, more particularly; a polypeptide starting at position 1658 and ending at position 1711, for expression of NS4a antigen, and more particularly, a polypeptide starting at position 1712 and ending in the region between positions 1743 and 1972 (for instance 1712-1743, 1712-1764, 1712-1782, 1712-1972, 1712-1782, 1712-1902), for expression of NS4b antigen or parts thereof.

Any other HCV vector construction known in the art may also be used for the recombinant polypeptides of the present invention.

Also any of the known purification methods for recombinant proteins may be used for the production of the recombinant polypeptides of the present invention, particularly the HCV recombinant polypeptide purification methods as disclosed in PCT/EP 95/03031 in name of Innogenetics N.V.

The term "vector" may comprise a plasmid, a cosmid, a phage, or a virus or

20

25

30

a transgenic animal. Particularly useful for vaccine development may be BCG or adenoviral vectors, as well as avipox recombinant viruses.

The present invention also relates to a method for the production of a recombinant polypeptide as defined above, comprising:

- transformation of an appropriate cellular host with a recombinant vector, in which a polynucleic acid or a part thereof according to as defined above has been inserted under the control of appropriate regulatory elements,
 - culturing said transformed cellular host under conditions enabling the expression of said insert, and,
- 10 harvesting said polypeptide.

The term 'recombinantly expressed' used within the context of the present invention refers to the fact that the proteins of the present invention are produced by recombinant expression methods be it in prokaryotes, or lower or higher eukaryotes as discussed in detail below.

The term 'lower eukaryote' refers to host cells such as yeast, fungi and the like. Lower eukaryotes are generally (but not necessarily) unicellular. Preferred lower Saccharomyces, species within particularly yeasts, eukaryotes are Schizosaccharomyces, Kluveromyces, Pichia (e.g. Pichia pastoris), Hansenula (e.g. Schizosaccharomyces, Schwaniomyces, Yarowia, polymorpha), Hanse<u>nula</u> Zygosaccharomyces and the like. Saccharomyces cerevisiae, S. carlsbergensis and K. lactis are the most commonly used yeast hosts, and are convenient fungal hosts.

The term 'prokaryotes' refers to hosts such as <u>E.coli</u>, <u>Lactobacillus</u>, <u>Lactococcus</u>, <u>Salmonella</u>, <u>Streptococcus</u>, <u>Bacillus subtilis</u> or <u>Streptomyces</u>. Also these hosts are contemplated within the present invention.

The term 'higher eukaryote' refers to host cells derived from higher animals, such as mammals, reptiles, insects, and the like. Presently preferred higher eukaryote host cells are derived from Chinese hamster (e.g. CHO), monkey (e.g. COS and Vero cells), baby hamster kidney (BHK), pig kidney (PK15), rabbit kidney 13 cells (RK13), the human osteosarcoma cell line 143 B, the human cell line HeLa and human hepatoma cell lines like Hep G2, and insect cell lines (e.g. Spodoptera frugiperda). The host cells may be provided in suspension or flask cultures, tissue cultures, organ cultures and the like. Alternatively the host cells may also be transgenic animals.

The term 'recombinant polynucleotide or nucleic acid' intends a polynucleotide

10

15

20

25

30

or nucleic acid of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than that to which it is linked in nature, or (3) does not occur in nature.

The term 'recombinant host cells', 'host cells', 'cells', 'cell lines', 'cell cultures', and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be or have been, used as recipients for a recombinant vector or other transfer polynucleotide, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

The term 'replicon' is any genetic element, e.g., a plasmid, a chromosome, a virus, a cosmid, etc., that behaves as an autonomous unit of polynucleotide replication within a cell; i.e., capable of replication under its own control.

The term 'vector' is a replicon further comprising sequences providing replication and/or expression of a desired open reading frame.

The term 'control sequence' refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, splicing sites and terminators; in eukaryotes, generally, such control sequences include promoters, splicing sites, terminators and, in some instances, enhancers. The term 'control sequences' is intended to include, at a minimum, all components whose presence is necessary for expression, and may also include additional components whose presence is advantageous, for example, leader sequences which govern secretion.

The term 'promoter' is a nucleotide sequence which is comprised of consensus sequences which allow the binding of RNA polymerase to the DNA template in a manner such that mRNA production initiates at the normal transcription initiation site for the adjacent structural gene.

The expression 'operably linked' refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their

10

15

20

25

30

intended manner. A control sequence 'operably linked' to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

The segment of the HCV cDNA encoding the desired sequence inserted into the vector sequence may be attached to a signal sequence. Said signal sequence may be that from a non-HCV source, e.g. the IgG or tissue plasminogen activator (tpa) leader sequence for expression in mammalian cells, or the α -mating factor sequence for expression into yeast cells, but particularly preferred constructs according to the present invention contain signal sequences appearing in the HCV genome before the respective start points of the proteins.

A variety of vectors may be used to obtain recombinant expression of HCV single or specific oligomeric envelope proteins of the present invention. Lower eukaryotes such as yeasts and glycosylation mutant strains are typically transformed with plasmids, or are transformed with a recombinant virus. The vectors may replicate within the host independently, or may integrate into the host cell genome.

Higher eukaryotes may be transformed with vectors, or may be infected with a recombinant virus, for example a recombinant vaccinia virus. Techniques and vectors for the insertion of foreign DNA into vaccinia virus are well known in the art, and utilize, for example homologous recombination. A wide variety of viral promoter sequences, possibly terminator sequences and poly(A)-addition sequences, possibly enhancer sequences and possibly amplification sequences, all required for the mammalian expression, are available in the art. Vaccinia is particularly preferred since vaccinia halts the expression of host cell proteins. Vaccinia is also very much preferred since it allows the expression of f.i. E1 and E2 proteins of HCV in cells or individuals which are immunized with the live recombinant vaccinia virus. For vaccination of humans the avipox and Ankara Modified Virus (AMV) are particularly useful vectors.

Also known are insect expression transfer vectors derived from baculovirus Autographa californica nuclear polyhedrosis virus (AcNPV), which is a helper-independent viral expression vector. Expression vectors derived from this system usually use the strong viral polyhedrin gene promoter to drive the expression of heterologous genes. Different vectors as well as methods for the introduction of heterologous DNA into the desired site of baculovirus are available to the man skilled

15

20

in the art for baculovirus expression. Also different signals for posttranslational modification recognized by insect cells are known in the art.

The present invention also relates to a host cell transformed with a recombinant vector as defined above.

The present invention also relates to a method for detecting antibodies to HCV present in a biological sample, comprising:

- (i) contacting the biological sample to be analysed for the presence of HCV with a polypeptide as defined above,
- (ii) detecting the immunological complex formed between said antibodies and said polypeptide.

The present invention also relates to a method for HCV typing, comprising: (i) contacting the biological sample to be analysed for the presence of HCV with a

- polypeptide as defined above,
- (ii) detecting the immunological complex formed between said antibodies and said polypeptide.

The present invention also relates to a diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one polypeptide as defined above, with said polypeptide being preferably bound to a solid support.

The present invention also relates to a diagnostic kit for HCV typing, said kit comprising at least one polypeptide as defined above, with said polypeptide being preferably bound to a solid support.

The present invention also relates to diagnostic kit according as defined above, said kit comprising a range of said polypeptides which are attached to specific locations on a solid substrate.

The present invention also relates to a diagnostic kit as defined above, wherein said solid support is a membrane strip and said polypeptides are coupled to the membrane in the form of parallel lines.

The immunoassay methods according to the present invention may utilize antigens from the different domains of the new and unique polypeptide sequences of the present invention that maintain linear (in case of peptides) and conformational epitopes (in case of polypeptides) recognized by antibodies in the sera from individuals infected with HCV. It is within the scope of the invention to use for instance single or specific oligomeric antigens, dimeric antigens, as well as

25

10

15

20

25

30

combinations of single or specific oligomeric antigens. The HCVantigens of the present invention may be employed in virtually any assay format that employs a known antigen to detect antibodies. Of course, a format that denatures the HCV conformational epitope should be avoided or adapted. A common feature of all of these assays is that the antigen is contacted with the body component suspected of containing HCV antibodies under conditions that permit the antigen to bind to any such antibody present in the component. Such conditions will typically be physiologic temperature, pH and ionic strength using an excess of antigen. The incubation of the antigen with the specimen is followed by detection of immune complexes comprised of the antigen.

Design of the immunoassays is subject to a great deal of variation, and many formats are known in the art. Protocols may, for example, use solid supports, or immunoprecipitation. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, enzymatic, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the immune complex are also known; examples of which are assays which utilize biotin and avidin or streptavidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

The immunoassay may be, without limitation, in a heterogeneous or in a homogeneous format, and of a standard or competitive type. In a heterogeneous format, the polypeptide is typically bound to a solid matrix or support to facilitate separation of the sample from the polypeptide after incubation. Examples of solid supports that can be used are nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates, polyvinylidine fluoride (known as ImmunolonTM), diazotized paper, nylon membranes, activated beads, and Protein A beads. For example, Dynatech ImmunolonTM 1 or ImmunlonTM 2 microtiter plates or 0.25 inch polystyrene beads (Precision Plastic Ball) can be used in the heterogeneous format. The solid support containing the antigenic polypeptides is typically washed after separating it from the test sample, and prior to detection of bound antibodies. Both standard and competitive formats are know in the art.

In a homogeneous format, the test sample is incubated with the combination of antigens in solution. For example, it may be under conditions that will precipitate

any antigen-antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art.

In a standard format, the amount of HCV antibodies in the antibody-antigen complexes is directly monitored. This may be accomplished by determining whether labeled anti-xenogeneic (e.g. anti-human) antibodies which recognize an epitope on anti-HCV antibodies will bind due to complex formation. In a competitive format, the amount of HCV antibodies in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labeled antibody (or other competing ligand) in the complex.

10

5

Complexes formed comprising anti-HCV antibody (or in the case of competitive assays, the amount of competing antibody) are detected by any of a number of known techniques, depending on the format. For example, unlabeled HCV antibodies in the complex may be detected using a conjugate of anti-xenogeneic lg complexed with a label (e.g. an enzyme label).

15

In an immunoprecipitation or agglutination assay format the reaction between the HCV antigens and the antibody forms a network that precipitates from the solution or suspension and forms a visible layer or film of precipitate. If no anti-HCV antibody is present in the test specimen, no visible precipitate is formed.

20

There currently exist three specific types of particle agglutination (PA) assays. These assays are used for the detection of antibodies to various antigens when coated to a support. One type of this assay is the hemagglutination assay using red blood cells (RBCs) that are sensitized by passively adsorbing antigen (or antibody) to the RBC. The addition of specific antigen antibodies present in the body component, if any, causes the RBCs coated with the purified antigen to agglutinate.

25

To eliminate potential non-specific reactions in the hemagglutination assay, two artificial carriers may be used instead of RBC in the PA. The most common of these are latex particles. However, gelatin particles may also be used. The assays utilizing either of these carriers are based on passive agglutination of the particles coated with purified antigens.

30

The HCV antigens of the present invention comprised of conformational epitopes will typically be packaged in the form of a kit for use in these immunoassays. The kit will normally contain in separate containers the native HCV antigen, control antibody formulations (positive and/or negative), labeled antibody

when the assay format requires the same and signal generating reagents (e.g. enzyme substrate) if the label does not generate a signal directly. The native HCV antigen may be already bound to a solid matrix or separate with reagents for binding it to the matrix. Instructions (e.g. written, tape, CD-ROM, etc.) for carrying out the assay usually will be included in the kit.

Immunoassays that utilize the native HCV antigen are useful in screening blood for the preparation of a supply from which potentially infective HCV is lacking. The method for the preparation of the blood supply comprises the following steps. Reacting a body component, preferably blood or a blood component, from the individual donating blood with HCV polypeptides of the present invention to allow an immunological reaction between HCV antibodies, if any, and the HCV antigen. Detecting whether anti-HCV antibody - HCV antigen complexes are formed as a result of the reacting. Blood contributed to the blood supply is from donors that do not exhibit antibodies to the native HCV antigens.

15

20

25

30

5

10

In cases of a positive reactivity to the HCV antigen, it is preferable to repeat the immunoassay to lessen the possibility of false positives. For example, in the large scale screening of blood for the production of blood products (e.g. blood transfusion, plasma, Factor VIII, immunoglobulin, etc.) 'screening' tests are typically formatted to increase sensitivity (to insure no contaminated blood passes) at the expense of specificity; i.e. the false-positive rate is increased. Thus, it is typical to only defer for further testing those donors who are 'repeatedly reactive'; i.e. positive in two or more runs of the immunoassay on the donated sample. However, for confirmation of HCV-positivity, the 'confirmation' tests are typically formatted to increase specificity (to insure that no false-positive samples are confirmed) at the expense of sensitivity.

The solid phase selected can include polymeric or glass beads, nitrocellulose,

microparticles, microwells of a reaction tray, test tubes and magnetic beads. The signal generating compound can include an enzyme, a luminescent compound, a chromogen, a radioactive element and a chemiluminescent compound. Examples of enzymes include alkaline phosphatase, horseradish peroxidase and beta-galactosidase. Examples of enhancer compounds include biotin, anti-biotin and avidin. Examples of enhancer compounds binding members include biotin, anti-biotin and avidin. In order to block the effects of rheumatoid factor-like substances, the

10

15

20

test sample is subjected to conditions sufficient to block the effect of rheumatoid factor-like substances. These conditions comprise contacting the test sample with a quantity of anti-human IgG to form a mixture, and incubating the mixture for a time and under conditions sufficient to form a reaction mixture product substantially free of rheumatoid factor-like substance.

The present invention particularly relates to an immunoassay format in which the polypeptides (or peptides) of the invention are coupled to a membrane in the form of parallel lines. This assay format is particularly advantageous for HCV typing purposes.

The present invention also relates to a pharmaceutical composition comprising at least one (recombinant) polypeptides as defined above and a suitable excipient, diluent or carrier.

The present invention also relates to a method of preventing HCV infection, comprising administering the pharmaceutical composition as defined above to a mammal in effective amount to stimulate the production of protective antibody or protective T-cell response.

The present invention relates to the use of a composition as defined above in a method for preventing HCV infection.

The present invention further relates to a vaccine for immunizing a mammal against HCV infection, comprising at least one (recombinant) polypeptide as defined above, in a pharmaceutically acceptable carrier.

The term 'immunogenic' refers to the ability of a substance to cause a humoral and/or cellular response, whether alone or when linked to a carrier, in the presence or absence of an adjuvant. 'Neutralization' refers to an immune response that blocks the infectivity, either partially or fully, of an infectious agent. A 'vaccine' is an immunogenic composition capable of eliciting protection against HCV, whether partial or complete. A vaccine may also be useful for treatment of an individual, in which case it is called a therapeutic vaccine.

The term 'therapeutic' refers to a composition capable of treating HCV infection. The term 'effective amount' refers to an amount of epitope-bearing polypeptide sufficient to induce an immunogenic response in the individual to which it is administered, or to otherwise detectably immunoreact in its intended system (e.g., immunoassay). Preferably, the effective amount is sufficient to effect

25

10

15

20

25

30

treatment, as defined above. The exact amount necessary will vary according to the application. For vaccine applications or for the generation of polyclonal antiserum / antibodies, for example, the effective amount may vary depending on the species, age, and general condition of the individual, the severity of the condition being treated, the particular polypeptide selected and its mode of administration, etc. It is also believed that effective amounts will be found within a relatively large, non-critical range. An appropriate effective amount can be readily determined using only routine experimentation. Preferred ranges of proteins for prophylaxis of HCV disease are 0.01 to 100 μ g/dose, preferably 0.1 to 50 μ g/dose. Several doses may be needed per individual in order to achieve a sufficient immune response and subsequent protection against HCV disease.

The present invention also relates to a vaccine as defined above, comprising at least one (recombinant) polypeptide as defined above, with said polypeptide being unique for at least one of the subtypes or types as defined above.

Said vaccine compositions may include prophylactic as well as therapeutic vaccine compositions.

Pharmaceutically acceptable carriers include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers; and inactive virus particles. Such carriers are well known to those of ordinary skill in the art.

Preferred adjuvants to enhance effectiveness of the composition include, but are not limited to: aluminim hydroxide (alum), N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP) as found in U.S. Patent No. 4,606,918, N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE) and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate, and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. Any of the 3 components MPL, TDM or CWS may also be used alone or combined 2 by 2. Additionally, adjuvants such as Stimulon (Cambridge Bioscience, Worcester, MA)

10

15

20

25

30

Immunogenic compositions used as vaccines comprise a 'sufficient amount' or 'an immunologically effective amount' of the proteins of the present invention, as well as any other of the above mentioned components, as needed. 'Immunologically effective amount', means that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment, as defined above. This amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated (e.g. nonhuman primate, primate, etc.), the capacity of the individual's immune system to synthesize antibodies, the degree of protection desired, the formulation of the vaccine, the treating doctor's assessment of the medical situation, the strain of infecting HCV, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials. Usually, the amount will vary from 0.01 to 1000 μ g/dose, more particularly from 0.1 to 100 μ g/dose.

The proteins of the invention may also serve as vaccine carriers to present homologous (e.g. T cell epitopes or B cell epitopes fromfor istance the core,E1, E2, NS2, NS3, NS4 or NS5 regions) or heterologous (non-HCV) haptens, in the same manner as Hepatitis B surface antigen (see European Patent Application 174,444). In this use, envelope proteins provide an immunogenic carrier capable of stimulating an immune response to haptens or antigens conjugated to the aggregate. The antigen may be conjugated either by conventional chemical methods, or may be cloned into the gene encoding E1 and/or E2 at a location corresponding to a hydrophilic region

10

15

20

25

30

of the protein. Such hydrophylic regions include the V1 region (encompassing amino acid positions 191 to 202), the V2 region (encompassing amino acid positions 213 to 223), the V3 region (encompassing amino acid positions 230 to 242), the V4 region (encompassing amino acid positions 230 to 242), the V5 region (encompassing amino acid positions 294 to 303) and the V6 region (encompassing amino acid positions 329 to 336). Another useful location for insertion of haptens is the hydrophobic region (encompassing approximately amino acid positions 264 to 293). It is shown in the present invention that this region can be deleted without affecting the reactivity of the deleted E1 protein with antisera. Therefore, haptens may be inserted at the site of the deletion.

The immunogenic compositions are conventionally administered parenterally, typically by injection, for example, subcutaneously or intramuscularly. Additional formulations suitable for other methods of administration include oral formulations and suppositories. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

The administration of the immunogen(s) of the present invention may be for either a prophylactic or therapeutic purpose. When provided prophylactically, the immunogen(s) is provided in advance of any exposure to HCV or in advance of any symptom of any symptoms due to HCV infection. The prophylactic administration of the immunogen serves to prevent or attenuate any subsequent infection of HCV in a mammal. When provided therapeutically, the immunogen(s) is provided at (or shortly after) the onset of the infection or at the onset of any symptom of infection or disease caused by HCV. The therapeutic administration of the immunogen(s) serves to attenuate the infection or disease.

In addition to use as a vaccine, the compositions can be used to prepare antibodies to HCV (E1) proteins. The antibodies can be used directly as antiviral agents. To prepare antibodies, a host animal is immunized using the E1 proteins native to the virus particle bound to a carrier as described above for vaccines. The host serum or plasma is collected following an appropriate time interval to provide a composition comprising antibodies reactive with the (E1) protein of the virus particle. The gamma globulin fraction or the IgG antibodies can be obtained, for example, by use of saturated ammonium sulfate or DEAE Sephadex, or other

techniques known to those skilled in the art. The antibodies are substantially free of many of the adverse side effects which may be associated with other anti-viral agents such as drugs.

The present invention also relates particularly to a peptide corresponding to an amino acid sequence encoded by at least one of the HCV genomic sequences as defined above, with said peptide being unique to any of the HCV subtypes or types as defined in Table 5, and which contains at least one amino acid differing from any of the known HCV types or subtypes, or an analog thereof being substantially homologous and biologically equivalent.

10

5

The present invention relates particularly to a peptide comprising at least one unique epitope of the new sequences of the invention as represented in SEQ ID NO 1 to 106.

The present invention relates also particularly to a peptide comprising in its sequence a unique amino acid residue of the invention as defined above.

15

The present invention relates particularly to a peptide which is biotinylated as explained in WO 93/18054.

All the embodiments (immunoassay formats, vaccines, compositions, uses, etc.) illustrated for the polypeptides of the invention as above also relate to the peptides of the invention.

20

25

The present invention also relates to a method for detecting antibodies to HCV present in a biological sample, comprising:

- (i) contacting the biological sample to be analysed for the presence of HCV with a peptide as defined above,
- (ii) detecting the immunological ccomplex formed between said antibodies and said peptide.

The present invention also relates to a method for HCV typing, comprising: (i) contacting the biological sample to be analysed for the presence of HCV with a peptide as defined above.

(ii) detecting the immunological complex formed between said antibodies and saidpeptide.

The present invention also relates to a diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one peptide as defined above, with said peptide being preferably bound to a solid support.

10

15

20

25

30

The present invention also relates to a diagnostic kit for HCV typing, said kit comprising at least one peptide as defined above, with said peptide being preferably bound to a solid support.

The present invention also relates to a diagnostic kit as defined above, wherein said peptides are selected from the following:

- at least one NS4 peptide,
- at least one NS4 peptide and at least one Core peptide,
- at least one NS4 peptide and at least one Core peptide and at least one E1 peptide,
- at least one NS4 peptide and at least one E1 peptide.

The present invention also relates to a diagnostic kit as defined above, said kit comprising a range of said peptides which are attached to specific locations on a solid substrate.

The present invention also relates to a diagnostic kit as defined above, wherein said solid support is a membrane strip and said peptides are coupled to the membrane in the form of parallel lines.

The present invention also relates to a pharmaceutical composition comprising at least one as defined above and a suitable excipient, diluent or carrier.

the present invention also relates to a method of preventing HCV infection, comprising administering the pharmaceutical composition as defined above to a mammal in effective amount to stimulate the production of protective antibody or protective T-cell response.

The present invention also relates to the use of a composition as defined above in a method for preventing HCV infection.

The present invention also relates to a vaccine for immunizing a mammal against HCV infection, comprising at least one peptide as defined above, in a pharmaceutically acceptable carrier.

The present invention relates also to a vaccine as defined above, comprising at least one peptide as defined above, with said peptide being unique for at least one of the subtypes or types as defined in Table 5.

The present invention relates to an antibody raised upon immunization with at least one polypeptide or peptide as defined above, with said antibody being specifically reactive with any of said polypeptides or peptides, and with said antibody being preferably a monoclonal antibody.

The monoclonal antibodies of the invention can be produced by any hybridoma liable to be formed according to classical methods from splenic cells of an animal, particularly from a mouse or rat, immunized against the HCV polypeptides according to the invention as defined above on the one hand, and of cells of a myeloma cell line on the other hand, and to be selected by the ability of the hybridoma to produce the monoclonal antibodies recognizing the polypeptides which has been initially used for the immunization of the animals.

The antibodies involved in the invention can be labelled by an appropriate label of the enzymatic, fluorescent, or radioactive type.

10

5

The monoclonal antibodies according to this preferred embodiment of the invention may be humanized versions of mouse monoclonal antibodies made by means of recombinant DNA technology, departing from parts of mouse and/or human genomic DNA sequences coding for H and L chains or from cDNA clones coding for H and L chains.

15

20

25

Alternatively the monoclonal antibodies according to this preferred embodiment of the invention may be human monoclonal antibodies. These antibodies according to the present embodiment of the invention can also be derived from human peripheral blood lymphocytes of patients infected with HCV type 1 subtype 1d, 1e, 1f or 1g, HCV type 2 subtype 2e, 2f, 2g, 2h, 2i, 2k or 2l; HCV type 3, subtype 3g; HCV type 4 subtype 4k, 4l or 4m; and/or HCV type 7 (subtypes 7a, 7c or 7d), 9, 10 or 11, or vaccinated against HCV. Such human monoclonal antibodies are prepared, for instance, by means of human peripheral blood lymphocytes (PBL) repopulation of severe combined immune deficiency (SCID) mice (for recent review, see Duchosal et al. 1992) or by screening Eppstein Barr-virustransformed lymphocytes of infected or vaccinated individuals for the presence of reactive B-cells by means of the antigens of the present invention.

The invention also relates to the use of the proteins of the invention, muteins thereof, or peptides derived therefrom for the selection of recombinant antibodies by the process of repertoire cloning (Persson et al., 1991).

30

Antibodies directed to peptides derived from a certain genotype may be used either for the detection of such HCV genotypes, or as therapeutic agents.

The present invention relates also to a method for detecting HCV antigens present in a biological sample, comprising:

15

20

25

30

- (i) contacting said biological sample with an antibody as defined above,
- (ii) detecting the immune complexes formed between said HCV antigens and said antibody.

The present invention relates also to a method for HCV typing, comprising:

- 5 (i) contacting said biological sample with an antibody as defined above,
 - (ii) detecting the immune complexes formed between said HCV antigens and said antibody.

The present invention relates also to a diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one antibody as defined above, with said antibody being preferably bound to a solid support.

The present invention relates also to a diagnostic kit for HCV typing, said kit comprising at least one antibody as defined above, with said antibody being preferably bound to a solid support.

The present invention relates also to a diagnostic kit as defined above, said kit comprising a range of said antibodies which are attached to specific locations on a solid substrate.

The present invention relates also to a pharmaceutical composition comprising at least one antibody as defined above and a suitable excipient, diluent or carrier.

The present invention relates also to a method of preventing or treating HCV infection, comprising administering the pharmaceutical composition as defined above to a mammal in effective amount.

The present invention relates also to the use of a composition as defined above in a method for preventing or treating HCV infection.

The genotype may also be detected by means of a type-specific antibody as defined above, which may also linked to any polynucleotide sequence that can afterwards be amplified by PCR to detect the immune complex formed (Immuno-PCR, Sano et al., 1992).

Any publications or patent applications referred to herein are incorporated by reference. The following examples illustrate aspects of the invention but are in no way intended to limit the scope thereof.

WO 96/13590

49

FIGURE LEGENDS

Figure Legends

Figure 1

Alignment of the nucleotide sequences of the Core/E1 region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

Figure 2

5

10

Alignment of the amino acid sequences of the Core/E1 region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

Figure 3

Nucleotide and amino acid sequences obtained from the new HCV isolates of the present invention (SEQ ID NO 1 to 106).

Figure 4

Alignment of the amino acid sequences of the Core/E1 region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

Figure 5

20

Alignment of the nucleotide sequences of the NS5b region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

Figure 6

Alignment of the amino acid sequences of the NS5b region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

5 Table 5

Overview of the new subtypes and types of the present invention and the regions sequenced. The subtypes between barckets have been replaced by the non-bracketed subtypes following the classification of Tokita et al. (1994).

Examples

10 Serum samples.

15

25

Serum samples from Cameroonian blood donors (CAM) were screened for HCV antibodies with Innotest HCV Ab III, and confirmed by INNO-LIA HCV III (Innogenetics, Antwerp, Belgium). Serum samples from patients with chronic hepatitis C infection were obtained from various centers in the Benelux countries (BNL), from France (FR), from Pakistan (PAK), from Egypt (EG), and from Vietnam (VN).

Samples from the Benelux, Cameroon, France and Vietnam were selected because of their aberrant reactivities (isolates CAM1078, FR2, FR1, VN4, VN12, VN13, NE98 and others (see Table 5)).

20 cPCR, LiPA, cloning and sequencing.

RNA isolation, cDNA synthesis, PCR, cloning, and LiPA genotyping using biotinylated 5' UR amplification products were performed as described (Stuyver et al., 1994c). The 5' UR, the Core/E1, and the NS5B PCR products were used for direct sequencing. The sequence of the universal 5' UR primers HCPr95, HCPr96, HCPr98, and HCPr29, were described previously (Stuyver et al. 1993b). The following primers were also described (Stuyver et al. 1994c): HCPr41, a sense

primer for the amplification of the Core region; HCPr52 and HCPr54 for amplification of the Core/E1 region; and HCPr206 and HCPr207 for amplication of a 340-bp NS5B region.

Serum samples BNL1, BNL2, BNL3, BNL4, BNL5, BNL6, BNL7, BNL8, BNL9, BNL10, BNL11, BNL12, CAM1078, FR2, FR16, FR4, FR13, VN13, VN4, VN12, FR1, 5 NE98, and FR19 were analyzed in the Core/E1 region by direct sequencing. Serum samples BNL1, BNL2, FR17, CAM1078, FR2, FR16, BNL3, FR4, BNL5, FR13, FR18, PAK64, BNL8, BNL12, EG81, VN13, VN4, VN12, FR1, NE98, FR14, FR15, and FR19 were also analyzed in the NS5B region by direct sequencing. Partial 5' UR, 10 Core, E1, and NS5B sequences were obtained. The length of the obtained sequences is sufficient to classify the obtained sequences into new types or subtypes, based on the phylogenetic distances to known sequences. The following sequences could be obtained (nucleotide sequences have odd-numbered SEQ ID NO., amino acid sequences have even-numbered SEQ ID NO.): SEQ ID NO 1, 3, 5, 7, 9, 11, 13, 15, 15 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103 and 105. The amino acid sequences deduced therefrom are given in SEQ ID NO 2, 4, 7, 9, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 20 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104 and 106. Table 5 gives an overview of these sequences.

	٥							2 9	/932-82/1(SEQ ID NO. / I)	157 ON CI 03311209 2002	/832-82/ [(SEG ID INO. / 5)	9	7932-8271(SEQ ID NO. 77) of	7932-8271(SEQ ID NO. 81)	9	7932-8271 (SEQ ID NO. 83)				1 (SEQ ID NO.	1 (SEQ ID NO.	1 (SEQ ID NO.	(SEQ ID NO.	ġ Z	7932-8271 (SEQ ID NO. 95)	7932-8271 (SEQ ID NO. 97)	7932-8266 (SEQ ID NO. 99)	٦.	7932-8271 (SEQ ID NO. 105)
	tion	478-925 (SEQ ID NO. 3)	478-925 (SEQ ID NO. 7)		(-238)-414 (SEQ ID NO. 59)		5)	478-957 (SEQ ID NO. 15)		(SEQ ID NO.	(SEQ ID NO.	478-833 (SEQ ID NO. 25)	75)		478-925 (SEQ ID NO. 29)	(SEQ ID	(SEQ ID NO.	(SEO ID	<u>Б</u> 80.	478-925 (SEQ ID NO. 39)						478-925 (SEQ ID NO. 51)			103)
	Nucleotide sequence position		1-310 (SEQ ID NO. 5)		1-223 (SEQ ID NO. 9)	1-950 (SEQ ID NO. 11)	(-15)-816 (SEQ ID NO. 6	1-310 (SEQ ID NO. 13)	1-957 (SEQ ID NO 17)		1-310 (SEQ ID NO. 21)		(-238)-957 (SEQ ID NO.		1-310 (SEQ ID NO. 27)							1-413 (SEQ ID NO. 45)	1-957 (SEQ ID NO. 43)	(SEO ID	(SEO ID	1-310 (SEO ID NO. 49)			(-238)-223 (SEQ ID NO. 103)
Table 5	Type Isolate	1d BNL1	1d BNL2	1d FR17	1e CAM1078	1f FR2	1g FR16	2e BNL3	2f FR4		2h BNL5				4k BNL7						F	8b)	7c(8a) VN4	74(9a) VN12	00(70) ER1	102 NE98			11a FR19

6 76/13370				53		PC1/EP95/04155						
2645-2757 (SEQ ID NO. 54) 2645-2757 (SEQ ID NO. 56)	(SEQ ID NO. (SEQ ID NO. (SEQ ID NO.	(SEQ 1D (SEQ 1D (SEQ 1D	(SEQ ID NO.	2645-2757 (SEQ ID NO. 78) (2645-2757 (SEQ ID NO. 80) (2645-2757 (SEQ ID NO. 82)	2645-2757 (SEQ ID NO. 84)	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2645-2757 (SEU ID NO. 98) 2645-2755 (SEQ ID NO. 100) 2645-2757 (SEQ ID NO. 102) 2645-2757 (SEQ ID NO. 106)					
sequence position 159-308 (SEQ ID NO. 4) 159-308 (SEQ ID NO. 8)	1-138 (SEQ ID NO. 60)	ID NO.	159-308 (SEQ ID NO. 20) 159-308 (SEQ ID NO. 24) 159-277 (SEQ ID NO. 26)	159-308 (SEQ ID NO. 30)	(SEQ ID NO.	308						
Amino acid 1-103 (SEQ ID NO. 2) 1-103 (SEQ ID NO. 6)	1-74 (SEQ ID NO. 10) 1-316 (SEQ ID NO. 12) 1-158 (SEQ ID NO. 66)	1-103 (SEQ ID NO. 14) 1-317 (SEQ ID NO. 18)	1-103 (SEQ ID NO. 22) 1-316 (SEQ ID NO. 76)	1-103 (SEQ ID NO. 28)		1-137 (SEQ ID NO. 46) 1-317 (SEQ ID NO. 44) 1-317 (SEQ ID NO. 48) 1-317 (SEQ ID NO. 42) 1-103 (SEQ ID NO. 50)	1-74 (SEQ ID NO. 104)					
Table 5-continued Type Isolate 1d BNL1 1d BNL2			2h BNL5 2i BNL6 2k FR13	21 FR18 3g PAK64 4k BNL7 4k BNL8		4m EG81 7a (8b)VN13 7c (8a)VN4 7d (9a)VN12 9a (7a)FR1 10a NE98 11a FR14						

Phylogenetic analysis.

Previously published sequences were taken from the EMBL/Genbank database. Alignments were created using the program HCVALIGN (Stuyver et al. 1994c). Sequences were presented in a sequential format to the Phylogeny Inference Package (PHYLIP) version 3.5c (public domain program freely available from the University of Washington, Seattle, USA). Distance matrices were produced by DNADIST using the Kimura 2-parameter setting and further analyzed in NEIGHBOR, using the neighborioning setting. The program DRAWTREE was used to create graphic outputs.

Identification of new subtypes

10

15

20

25

5

These analyses indicated the clustering of BNL1, BNL2, CAM 1078, FR2, FR16, and FR17 with type 1 isolates, yet neither of these sequences clustered together with any of the known type 1 subtypes 1a, 1b, or 1c. BNL1, BNL2, and FR17 clearly clustered together and could be assigned a new type 1 subtype 1d, while CAM1078 could be classified into another new subtype 1e, FR2 could be classified into another type 1 subtype 1f, and FR16 could be classified into yet another type 1 subtype 1g. Interestingly, all 3 type 1d isolates (BNL1, BNL2, and FR17) and 1g isolate FR16 were obtained from patients of Moroccan ethnic origin who resided in Europe.

Another group of isolates showed homology to other type 2 sequences, but none of the isolates BNL3, FR4, BNL4, BNL5, BNL6, FR13, or FR18 could be classified into one of the known type 2 subtypes 2a, 2b, 2c (Bukh et al., 1993), or 2d (Stuyver et al., 1994c). Based on the phylogenetic distances to other type 2 isolates and to other isolates of the group, each of these isolates could be classified into a new type 2 subtype. BNL3 was assigned subtype 2e, FR4 subtype 2f, BNL4 subtype 2g, BNL5 subtype 2h, and BNL6 could be classified into yet another type 2 subtype 2i. If the previously published isolate HN4 is classified as 2j, FR13 and FR18 may be classified into new type 2 subtypes 2k and 2l. However, the possibility that FR13 and FR18 could belong to subtypes 2g or 2i has not yet been ruled out. Definite classification can be obtained by determining the NS5B sequences of isolates BNL4 and BNL6, belonging to subtypes 2g and 2i, respectively.

30

Isolate PAK64 showed homology to type 3 sequences, but could not be classified into one of the known type 3 subtypes 3a to f. Based on the phylogenetic distances to other type 3 isolates, PAK64 could be classified into a new type 3

10

subtype. PAK64 was assigned subtype 3g. However, the possibility that PAK64 belongs to a known type 3 subtype can not be strictly ruled out since only one region of the genome has been sequenced. Definite classification can be obtained by determining the Core/E1 sequences of isolate PAK64 after amplification with primerHcPr52 and HcPr54.

Among the Benelux and Egyptian samples that were analyzed, some sequences clustered with the previously identified type 4 subtypes 4c and 4d. However, BNL7, BNL8, BNL9, BNL10, BNL11, BNL12, and EG81 clustered into new subtypes of type 4. Isolates BNL7, BNL8, BNL9, BNL10, and BNL11 clustered again separately from BNL12 and EG81 into a new subtype 4k. This subtype was the predominant subtype in the Benelux countries. BNL12 and EG81 also segregated into separate subtypes. BNL12 was assigned to another new subtype 4l and EG81 was assigned to yet another new subtype 4m.

10

15

Identification of new HCV major types

Isolates FR1, VN4, VN12, VN13, NE98, FR14, FR15, and FR19 did not cluster with any of the known 6 major types of HCV. VN4, VN12, and VN13 were very distantly related to genotype 6, but phylogenetic analysis indicated that these isolates should be assigned new major types. VN13, VN4 and VN12 were related at the subtype level and assigned type 7a, 7c, and 7d, respectively. FR1 was not related to any known isolate and was assigned genotype 9a. NE98 shows a distant relatedness to type 3 sequences, yet phylogenetic analysis suggested classification into a new major type 10a. Depending on international guidelines for assigning type and subtype levels, NE98 may also be classified into an additional type 3 subtype. FR14, FR15, and FR19 show a very distant relatedness to type 2 sequences, yet phylogenetic analysis indicated thes isolates to be classified into a new major type 11, all belonging to the same subtype designated 11a. Depending on international guidelines for assigning type and subtype levels, FR14, FR15, and FR19 may also be classified into an additional type 2 subtype.

REFERENCES

Barany F (1991). Genetic disease detection and DNA amplification using cloned thermostable ligase. Proc Natl Acad Sci USA 88: 189-193.

Bej A, Mahbubani M, Miller R, Di Cesare J, Haff L, Atlas R (1990) Mutiplex PCR amplification and immobilized capture probes for detection of bacterial pathogens and indicators in water. Mol Cell Probes 4:353-365.

Bukh J, Purcell R, Miller R (1992). Sequence analysis of the 5' noncoding region of hepatitis C virus. Proc Natl Acad Sci USA 89:4942-4946.

Bukh J, Purcell R, Miller R (1993). At least 12 genotypes of hepatitis C virus predicted by sequence analysis of the putative E1 gene of isolates collected worldwide. Proc. Natl. Acad. Sci. USA 90,8234-8238.

Cha T, Beal E, Irvine B, Kolberg J, Chien D, Kuo G, Urdea M (1992) At least five related, but distinct, hepatitis C viral genotypes exist. Proc Natl Acad Sci USA 89:7144-7148.

15 Chan S-W, Simmonds P, McOmish F, Yap P, Mitchell R, Dow B, Follett E (1991) Serological responses to infection with three different types of hepatitis C virus. Lancet 338:1991.

Chan S-W, McOmish F, Holmes E, Dow B, Peutherer J, Follett E, Yap P, Simmonds P (1992) Analysis of a new hepatitis C virus type and its phylogenetic relationship to existing variants. J Gen Virol 73:1131-1141.

Chomczynski P, Sacchi N (1987) Single step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. Anal Biochem 162:156-159.

Choo Q, Richman K, Han J, Berger K, Lee C, Dong C, Gallegos C, Coit D, Medina-

25

Selby A, Barr P, Weiner A, Bradley D, Kuo G, Houghton M (1991) Genetic organization and diversity of the hepatitis C virus. Proc Natl Acad Sci USA 88:2451-2455.

Compton J (1991). Nucleic acid sequence-based amplification. Nature, 350: 91-92.

Duchosal A, Eming S, Fisher P (1992) Immunization of hu-PBL-SCID mice and the resue of human monoclonal Fab fragments through combinatorial libraries. Nature 355:258-262.

Duck P (1990). Probe amplifier system based on chimeric cycling oligonucleotides. Biotechniques 9, 142-147.

Guatelli J, Whitfield K, Kwoh D, Barringer K, Richman D, Gengeras T (1990)

10 Isothermal, in vitro amplification of nucleic acids by a multienzyme reaction modeled after retroviral replication. Proc Natl Acad Sci USA 87: 1874-1878.

Hijikata M, Kato N, Ootsuyama Y, Nakagawa M, Shimotohmo K (1991) Gene mapping of the putative structural region of the hepatitis C virus genome by in vitro processing analysis. Proc Natl Acad Sci USA 88, 5547-5551.

Jacobs K, Rudersdorf R, Neill S, Dougherty J, Brown E, Fritsch E (1988) The thermal stability of oligonucleotide duplexes is sequence independent in tetraalkylammonium salt solutions: application to identifying recombinant DNA clones. Nucl Acids Res 16:4637-4650.

Kato N, Hijikata M, Ootsuyama Y, Nakagawa M, Ohkoshi S, Sugimura T, Shimotohno K (1990) Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis. Proc Natl Acad Sci USA 87:9524-9528.

Kwoh D, Davis G, Whitfield K, Chappelle H, Dimichele L, Gingeras T (1989). Transcription-based amplification system and detection of amplified human immunodeficiency virus type 1 with a bead-based sandwich hybridization format. Proc Natl Acad Sci USA, 86: 1173-1177.

SUBSTITUTE SHEET (RULE 26)

WO 96/13590 PCT/EP95/04155

Kwok S, Kellogg D, McKinney N, Spasic D, Goda L, Levenson C, Sinisky J, (1990). Effects of primer-template mismatches on the polymerase chain reaction: Human immunodeficiency views type 1 model studies. Nucl. Acids Res., 18: 999.

Landgren U, Kaiser R, Sanders J, Hood L (1988). A ligase-mediated gene detection technique. Science 241:1077-1080.

Lizardi P, Guerra C, Lomeli H, Tussie-Luna I, Kramer F (1988) Exponential amplification of recombinant RNA hybridization probes. Bio/Technology 6:1197-1202.

Lomeli H, Tyagi S, Printchard C, Lisardi P, Kramer F (1989) Quantitative assays based on the use of replicatable hybridization probes. Clin Chem 35: 1826-1831.

Machida A, Ohnuma H, Tsuda F, Munekata E, Tanaka T, Akahane Y, Okamoto H, Mishiro S (1992) Hepatology 16, 886-891.

Maniatis T, Fritsch E, Sambrook J (1982) Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

Mori S, Kato N, Yagyu A, Tanaka T, Ikeda Y, Petchclai B, Chiewsilp P, Kurimura T, Shimotohno K (1992) A new type of hepatitis C virus in patients in Thailand. Biochem Biophys Res Comm 183:334-342.

Okamoto H, Okada S, Sugiyama Y, Kurai K, lizuka H, Machida A, Miyakawa Y, Mayumi M (1991) Nucleotide sequence of the genomic RNA of hepatitis C virus isolated from a human carrier: comparison with reported isolates for conserved and divergent regions. J Gen Virol 72:2697-2704.

20

Okamoto H, Kurai K, Okada S, Yamamoto K, Lizuka H, Tanaka T, Fukuda S, Tsuda F, Mishiro S (1992) Full-length sequences of a hepatitis C virus genome having poor homology to reported isolates: comparative study of four distinct genotypes. Virology 188:331-341.

20

Persson M, Caothien R, Burton D (1991). Generation of diverse high-affinity human monoclonal antibodies by repertoire cloning. Proc Natl Acad Sci USA 89:2432-2436. Saiki R, Gelfand D, Stoffel S, Scharf S, Higuchi R, Horn G, Mullis K, Erlich H (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. Science 239:487-491.

Saiki R, Walsh P, Levenson C, Erlich H (1989) Genetic analysis of amplified DNA with immobilized sequence-specific oligonucleotide probes (1989) Proc Natl Acad Sci USA 86:6230-6234.

Sano T, Smith C, Cantor C (1992) Immuno-PCR: very sensitive antigen detection by means of specific antibody-DNA conjugates. Science 258:120-122.

Simmonds P, McOmsh F, Yap P, Chan S, Lin C, Dusheiko G, Saeed A, Holmes E (1993a), Sequence variability in the 5' non-coding region of hepatitis C virus : identification of a new virus type and restrictions on sequence diversity. J Gen Virology, 74:661-668.

Stuyver L, Rossau R, Wyseur A, Duhamel M, Vanderborght B, Van Heuverswyn H, Maertens G (1993b) Typing of hepatitis C virus (HCV) isolates and characterization of new (sub)types using a Line Probe Assay. J Gen Virology, 74: 1093-1102.

Tokita et al. (1994) Hepatitis C virus vraiants from Vietnam are classifiable into the seventh, eighth, and ninth major genetic groups. Proc. Natl. Acad. Sci, 91: 11022-11026.

Walker G, Little M, Nadeau J, Shank D (1992). Isothermal in vitro amplification of DNA by a restriction enzyme/DNA polymerase system. Proc Natl Acad Sci USA 89:392-396.

Wu D, Wallace B (1989). The ligation amplification reaction (LAR) - amplification of specific DNA sequences using sequential rounds of template-dependent ligation.

Genomics 4:560-569.

WO 96/13590

15

20

Miller P, Yano J, Yano E, Carroll C, Jayaram K, Ts'o P (1979) Nonionic nucleic acid analogues. Synthesis and characterization of dideoxyribonucleoside methylphosphonates. Biochemistry 18(23):5134-43.

Nielsen P, Egholm M, Berg R, Buchardt O (1991) Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. Science 254(5037):1497-500.

Nielsen P, Egholm M, Berg R, Buchardt O (1993) Sequence specific inhibition of DNA restriction enzyme cleavage by PNA. Nucleic-Acids-Res. 21(2):197-200.

Asseline U, Delarue M, Lancelot G, Toulme F, Thuong N (1984) Nucleic acid-binding molecules with high affinity and base sequence specificity: intercalating agents covalently linked to oligodeoxynucleotides. Proc. Natl. Acad. Sci. USA 81(11):3297-301.

Matsukura M, Shinozuka K, Zon G, Mitsuya H, Reitz M, Cohen J, Broder S (1987) Phosphorothioate analogs of oligodeoxynucleotides: inhibitors of replication and cytopathic effects of human immunodeficiency virus. Proc. Natl. Acad. Sci. USA 84(21):7706-10.

Maertens, G., Ducatteeuw, A., Stuyver, L., Vandeponseele, P., Venneman, A., Wyseur, A., Bosman, F., Heijtink, R. & de Martynoff, G. (1994) Low prevalence of anti-E1 antibodies reactive to recombinant type 1b E1 envelope protein in type 2, 3, and 4 sera, but high prevalence in subtypes 1a and 1b. In: Viral Hepatitis and Liver Disease, Proceedings of the International Symposium on Viral Hepatitis and Liver Disease (Eds. Nishioka, K., Suzuki, H., Mishiro, S., and Oda, T.), pp 314-316, Springer-Verlag Tokyo.

Simmonds, P., Rose, K.A., Graham, S., Chan, S.-W., McOmish, F., Dow, B.C., Follett,

E.A.C., Yap, P.L., & Marsden, H. (1993b) Mapping of serotype-specific, immunodominant epitopes in the NS4 region of hepatitis C virus (HCV): Use of type-specific peptides to serologically discriminate infections with HCV type 1, 2, and 3.

J. Clin. Microbiol. 31, 1493-1503.

Simmonds, P., Holmes, E.C., Cha, T.-A., Chan, S.-W., McOmish, F., Irvine, B., Beall, E., Yap, P.L., Kolberg, J., & Urdea, M.S. (1993c) *J. Gen. Virol.* **74**, 2391-2399.

Stuyver, L., Van Arnhem, W., Wyseur, A. & Maertens, G. (1994) Cloning and phylogenetic analysis of the Core, E2, and NS3/4 regions of hepatitis C virus type 5a. Biochem. Biophys. Res. Comm. 202, 1308-1314.

Simmonds, P., Alberti, A., Alter, H., Bonino, F., Bradley, D.W., Bréchot, C., Brouwer, J., Chan, S.-W., Chayama K., Chen, D.-S., Choo, Q.-L., Colombo, M., Cuypers, T., Date, T., Dusheiko, G., Esteban, J.I., Fay, O., Hadziyannis, S., Han, J., Hatzakis, A., Holmes, E.C., Hotta, H., Houghton, M., Irvine, B., Kohara, M., Kolberg, J.A., Kuo, G., Lau, J.Y.N., Lelie, P.N., Maertens, G., McOmish, F., Miyamura, T., Mizokami, M., Nomoto, A., Prince A.M., Reesink, H.W., Rice, C., Roggendorf, M., Schalm, S., Shikata, T., Shimotohno, K., Stuyver, L., Trépo, C., Weiner, A., Yap, P.L. & Urdea, M.S. (1994) A proposed system for the nomenclature of hepatitis C virus genotypes.
Hepatology 19, 1321-1324.

Stuyver, L., Van Arnhem, W., Wyseur, A., DeLeys, R. & Maertens, G. (1993a) Analysis of the putative E1 envelope and NS4a epitope regions of HCV type 3. Biochem. Biophys. Res. Comm. 192, 635-641.

Stuyver, L., Rossau, R., Wyseur, A., Duhamel, M., Vanderborght, B., Van Heuverswyn, H. & Maertens, G. (1993b) Typing of hepatitis C virus isolates and characterization of new subtypes using a line probe assay. J. Gen Virol. 74, 1093-1102.

Stuyver, L., Wyseur, A., Van Arnhem, W., Rossau, R., Delaporte, E., Dazza, M.-C., Van Doorn, L.-J., Kleter, B. & Maertens, G. (1994a) The use of a line probe assay as a tool to detect new types or subtypes of hepatitis C virus. In: Viral Hepatitis and Liver Disease, Proceedings of the International Symposium on Viral Hepatitis and Liver Disease (Eds. Nishioka, K., Suzuki, H., Mishiro, S., and Oda, T.), pp 317-319,

Springer-Verlag Tokyo.

Stuyver, L., Van Arnhem, W., Wyseur, A. & Maertens, G. (1994b) Cloning and Phylogenetic analysis of the Core, E2, and NS3/4 regions of the hepatitis C virus type 5a. Biochem. Biophys. Res. Comm. 202, 1308-1314.

Stuyver, L., Van Arnhem, W., Wyseur, A., Hernandez, F., Delaporte, E., & Maertens, G. (1994c) Classification of hepatitis C viruses based on phylogenetics analysis of the E1 and NS5B regions and identification of 5 new subtypes. Proc. Natl. Acad. Sci. USA 91.

Stuyver et al. (1995) Hepatitis C virus genotyping by means of 5'-UR/core line probe assays and molecular analysis of untypeable samples. Virus Reasearch (in press).

15

20

25

CLAIMS

- 1. An HCV polynucleic acid, having a nucleotide sequence which is unique to a theretofore unidentified HCV type or subtype which is different from HCV subtypes 1a, 1b, 1c, 2a, 2b, 2c, 2d, 3a, 3b, 3c, 3d, 3e, 3f, 4a, 4b, 4c, 4d, 4e, 4f, 4g, 4h, 4i, 4j, 5a or 6a, with said HCV subtypes being classified as in Table 3 by comparison of a part of the NS5 gene nucleotide sequence spanning positions 7932 to 8271, with said amino acid numbering being shown in Table 1, and with said polynucleic acid containing at least one nucleotide differing from said known HCV nucleotide sequences, or the complement thereof.
- A polynucleic acid according to claim 1, having a nucleotide sequence which is
 unique to at least one of HCV subtypes 1d, 1e, 1f, 1g, 2e, 2f, 2g, 2h, 2i, 2k, 2l, 3g,
 4k, 4l, 4m, 7a, 7c or 7d, with said HCV subtypes being classified as defined in claim
 1.
 - 3. A polynucleic acid according to claim 1, having a nucleotide sequence which is unique to at least one of HCV types 9, 10 or 11, with said HCV types being classified as defined in claim 1.
 - 4. A polynucleic acid according to any of claims 1 to 3 encoding an HCV polyprotein comprising in its amino acid sequence at least one of the following amino acid residues:
 - I15, C38, V44, A49, Q43, P49, Q55, A58, S60 or D60, E68 or V68, H70, A71 or Q71 or N71, D72, H81, H101, D106, S110, L130, I134, E135, L140, S148, T150 or E150, Q153, F155, D157, G160, E165, I169, F181, L186, T190, T192 or I192 or H192, I193, A195, S196, R197 or N197 or K197, Q199 or D199 or H199 or N199, F200 or T200, A208, I213, M216 or S216, N217 or S217 or G217 or K217, T218, I219, A222, Y223, I230, W231 or L231, S232 or H232 or A232, Q233, E235 or L235, F236 or T236, F237, L240 or M240, A242, N244, N249, I250 or K250 or R250, A252 or C252, A254, I255 or V255, D256 or M256, E257, E260 or K260, R261, V268, S272 or R272, I285, G290 or F290, A291, A293 or L293 or W293, T294 or A294, S295 or H295, K296 or E296, Y297 or M297, I299 or Y299, I300,

10

30

S301, P316, S2646, A2648, G2649, A2650, V2652, Q2653, H2656 or L2656, D2657, F2659, K2663 or Q2663, A2667 or V1667, D2677, L2681, M2686 or Q2686 or E2686, A2692 or K2692, H2697, I2707, L2708 or Y2708, A2709, A2719 or M2719, F2727, T2728 or D2728, E2729, F2730 or Y2730, I2741, I2745, V2746 or E2746 or L2746 or K2746, A2748, S2749 or P2749, R2750, E2751, D2752 or N2752 or S2752 or T2752 or V2752 or I2752 or Q2752, S2753 or D2753 or G2753, D2754, A2755, L2756 or Q2756, R2757,

with said notation being composed of a letter representing the amino acid residue by its one-letter code, and a number representing the amino acid numbering as shown in Table 1,

or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.

5. A polynucleic acid according to any of claims 1 to 4, with said polynucleic acid
 encoding a HCV polyprotein comprising in its amino acid sequence at least one amino acid sequence chosen from the following list:

ARQSDGRSWAQ or ARRSEGRSWAQ as for subtype 1d	(SEQ ID NO 107 and 108)
ERRPEGRSWAQ as for subtype 1e	(SEQ ID NO 109)
ARRPEGRSWAQ as for subtype 1f	(SEQ ID NO 110)
DRRTTGKSWGR as for subtype 2k	(SEQ ID NO 111)
DRRATGRSWGR as for subtype 2e	(SEQ ID NO 112)
DRRATGKSWGR as for subtype 2f	(SEQ ID NO 113)
VRQPTGRSWGQ as for type 9	(SEQ ID NO 114)
VRHQTGRTWAQ as for subtype 7a and 7c	(SEQ ID NO 115)
VRQNQGRTWAQ as for subtype 7d	(SEQ ID NO 116)
	(SEQ ID NO 117)
	(SEQ ID NO 118 and
119)	(ord in 140) to and
HEVRNASGVYHV or HEVRNASGVYHL as for subtype 1d	(SEQ ID NO 120 and 121)
	ERRPEGRSWAQ as for subtype 1e ARRPEGRSWAQ as for subtype 1f DRRTTGKSWGR as for subtype 2k DRRATGRSWGR as for subtype 2e DRRATGKSWGR as for subtype 2f VRQPTGRSWGQ as for type 9 VRHQTGRTWAQ as for subtype 7a and 7c VRQNQGRTWAQ as for subtype 7d ARRTEGRSWAQ as for type 10 VRRTTGRXXXXX or VRRTTGRTWAQ as for type 11

YEVHSTTDGYHV as for subtype 1f (SEQ ID NO 122)

VEVKNTSQAYMA as for subtype 2e (SEQ ID NO 123)

IQVKNNSHFYMA as for subtype 2f (SEQ ID NO 124)

	VQVKNTSTMYMA as for subtype 2g	(SEQ ID NO 125)
	VQVKNTSHSYMV as for subtype 2h	(SEQ ID NO 126)
	VQVANRSGSYMV as for subtype 2i	(SEQ ID NO 127)
	VEIKNTXNTYVL or VEIKNTSNTYVL as for so	ubtype 2k (SEQ ID NO 128 and 129)
5	INYRNVSGIYYV or INYRNTSGIYHV or INYH	
	subtype 4k	(SEQ ID NO 130, 131, 132 or 133)
	QHYRNVSGIYHV as for subtype 4I	(SEQ ID NO 134)
	IQVKNASGIYHL as for type 9	(SEQ ID NO 135)
	AHYTNKSGLYHL as for subtype 7c	(SEQ ID NO 136)
10	LNYANKSGLYHL as for subtype 7d	(SEQ ID NO 137)
	LEYRNASGLYMV as for type 10	(SEQ ID NO 138)
	IYEMDGMIMHY or IYEMSGMILHA as for sub	otype 1d (SEQ ID NO 139 and 140)
	VYEAKDIILHT as for subtype 1f	(SEQ ID NO 141)
	VWQLXDAVLHV as for subtype 2e	(SEQ ID NO 142)
15	VWQLRDAVLHV as for subtype 2f	(SEQ ID NO 143)
	IWQMQGAVLHV as for subtype 2g	(SEQ ID NO 144)
	VWQLKDAVLHV as for subtype 2h	(SEQ ID NO 145)
	VWQLEEAVLHV as for subtype 2i	(SEQ ID NO 146)
	TWQLXXAVLHV as for subtype 2k	(SEQ ID NO 147)
20	VYEADHHILHL or VYEADHHILAL or VFEADI	HILHL as for subtupe 4k
	(SEQ	ID NO 148, 149 and 150)
	VYESDHHILHL as for subtype 4l	(SEQ ID NO 151)
	VFEAETMILHL as for type 9	(SEQ ID NO 152)
	VYEAETLILHL as for subtype 7c	(SEQ ID NO 153)
25	VYEANGMILHL as for subtype 7d	(SEQ ID NO 154)
	VYEAGDIILHL as for type 10	(SEQ ID NO 155)
	VREDNHLRCWMAL or VRENNSSRCWMAL a	as for subtype 1d
		(SEQ ID NO 156 and 157)
	IREGNISRCWVPL as for subtype 1f	(SEQ ID NO 158)
30	ENSSGRFHCWIPI as for subtype 2e	(SEQ ID NO 159)
	ERSGNRTFCWTAV as for subtype 2f	(SEQ ID NO 160)
	ELQGNKSRCWIPV as for subtype 2g	(SEQ ID NO 162)
	ERHQNQSRCWIPV as for subtype 2h	(SEQ ID NO 163)

	~ ,	
	EWKDNTSRCWIPV as for subtype 2i	(SEQ ID NO 164)
	EREGNSSRCWIPV as for subtype 2k	(SEQ ID NO 165)
	VREGNOSRCWVAL or VRTGNOSRCWVAL	or VRVGNQSSCWVAL or
	VRVGNQSRCWVAL or VKEGNHSRCWVAL as for sul	otype 4k
5	(SEQ ID NO 1	66, 167, 168 or 169)
	VKTGNTSRCWVAL as for subtype 4I	(SEQ ID NO 170)
	IKAGNESRCWLPV as for type 9	(SEQ ID NO 171)
	VKEGNQSRCWVQA as for subtype 7c	(SEQ ID NO 172)
	VKXXNLTKCWLSA as for subtype 7d	(SEQ ID NO 173)
10	VRSGNTSRCWIPV as for type 10	(SEQ ID NO 174)
	VKNASVPTAA or VKDANVPTAA as for subtype 1d	(SEQ ID NO 175 and
	176)	
	ARIANAPIDE as for subtype 1f	(SEQ ID NO 177)
	VSKPGALTKG as for subtype 2e	(SEQ ID NO 178)
15	VSRPGALTRG as for subtype 2f	(SEQ ID NO 179)
	VNQPGALTRG as for subtype 2g	(SEQ ID NO 180)
	VSQPGALTRG as for subtype 2h	(SEQ ID NO 181)
	VSQPGALTKG as for subtype 2i	(SEQ ID NO 182)
	VSRPGALTEG as for subtype 2k	(SEQ ID NO 183)
20	APYIGAPLES or APYTAAPLES as for subtype 4k	(SEQ ID NO 184 and 185)
	APILSAPLMS as for subtype 4I	(SEQ ID NO 186)
	VPNSSVPIHG as for type 9	(SEQ ID NO 187)
	VPNASTPVTG as for subtype 7c	(SEQ ID NO 188)
	VQNASVSIRG as for subtype 7d	(SEQ ID NO 189)
25	VKSPCAATAS as for type 10	(SEQ ID NO 190)
	SPRMHHTTQE or SPRLYHTTQE as for subtype 1d	(SEQ ID NO 191 and 192)
	TSRRHWTVQD as for subtype 1f	(SEQ ID NO 193)
	APKRHYFVQE as for subtype 2e	(SEQ ID NO 194)
	SPQYHTFVQE as for subtype 2f	(SEQ ID NO 195)
30	SPQHHNFSQD as for subtype 2g	(SEQ ID NO 196)
	SPQHHIFVQD as for subtype 2h	(SEQ ID NO 197)
	SPEHHHFVQD as for subtype 2k	(SEQ ID NO 198)
	RPRRHWTTQD or RPRRHWTAQD or QPRRHWTTQD or	RPRRHWTTQE as for
	SUBSTITUTE SHEET (RULE 26)	

PCT/EP95/04155

WO 96/13590

5

20

25

68

Subtype 4k (SEQ ID NO 199, 200, 201 or 202)

QPRRHWTVQD as for subtype 4I (SEQ ID NO 203)

RPKYHQVTQD as for type 9 (SEQ ID NO 204)

RPRMHQVVQE as for subtype 7c (SEQ ID NO 205)

RPRMYEIAQD as for subtype 7d (SEQ ID NO 206)

RHRQHWTVQD as for type 10 (SEQ ID NO 207)

or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one nucleotide

6. A polynucleic acid according to any of claims 1 to 5 having a sequence selected from any of SEQ ID NO 1 to 105, or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.

differing from known HCV nucleotide sequences, or the complement thereof.

- 7. A polynucleic acid according to any of claims 1 to 6, which codes for the 5' UR, the Core/E1, the NS4 or the NS5B region or a part thereof.
 - 8. A polynucleic acid according to any of claims 1 to 7 which is a cDNA sequence.
 - 9. An oligonucleotide primer comprising part of a polynucleic acid according to any of claims 1 to 8, with said primer being able to act as primer for specifically amplifying the nucleic acid of a certain isolate belonging to the genotype from which the primer is derived.
 - 10. An oligonucleotide probe comprising part of a polynucleic acid according to any of claims 1 to 8, with said probe being able to act as a hybridization probe for specific detection and/or classification into types and/or subtypes of a HCV nucleic acid containing said nucleotide sequence, with said probe being possibly labelled or attached to a solid substrate.
 - 11. A diagnostic kit for use in determing the genotype of HCV, said kit comprising a

primer according to claim 9.

- 12. A diagnostic kit for use in determining the genotype of HCV, said kit comprising a probe according to claim 10.
- 13. A diagnostic kit according to claim 12, wherein said probe(s) is(are) attached to a solid substrate.
- 14. A diagnostic kit according to claim 13, wherein a range of said probes are attached to specific locations on a solid substrate.
- 15. A diagnostic kit according to claim 14, wherein said solid support is a membrane strip and said probes are coupled to the membrane in the form of parallel lines.
- 16. A method for the detection of HCV nucleic acids present in a biological sample, comprising:
 - (i) possibly extracting sample nucleic acid,
 - (ii) amplifying the nucleic acid with at least one primer according to claim 9,
 - (iii) detecting the amplified nucleic acids.
- 15 17. A method for the detection of HCV nucleic acids present in a biological sample, comprising:
 - (i) possibly extracting sample nucleic acid,
 - (ii) possibly amplifying the nucleic acid with at least one primer according to claim 9, or with a universal HCV primer,
- 20 (iii) hybridizing the nucleic acids of the biological sample, possibly under denatured conditions, at appropriate conditions with one or more probes according to claim 10, with said probes being possibly attached to a solid substrate,
 - (iv) possibly washing at appropriate conditions,
- 25 (v) detecting the hybrids formed.
 - 18. A method for detecting the presence of one or more HCV genotypes present in SUBSTITUTE SHEET (RULE 26)

a biological sample, comprising:

- (i) possibly extracting sample nucleic acid,
- (ii) specifically amplifying the nucleic acid with at least one primer according to claim 9,
- 5 (iii) detecting said amplified nucleic acids,
 - (iv) inferring the presence of one or more genotypes of HCV present from the observed pattern of amplified fragments.
 - 19. A method for detecting the presence of one or more HCV genotypes present in a biological sample, comprising:
- 10 (i) possibly extracting sample nucleic acid,
 - (ii) possibly amplifying the nucleic acid with at least one primer according to claim 9 or with a universal HCV primer,
 - (iii) hybridizing the nucleic acids of the biological sample, possibly under denatured conditions, at appropriate conditions with one or more probes according to claim 10, with said probes being possibly attached to a solid substrate,
 - (iv) possibly washing at appropriate conditions,
 - (v) detecting the hybrids formed,
- (vi) inferring the presence of one or more HCV genotypes present from the observed hybridization pattern.
 - 20. A method according to claim 19, wherein said probes are further characterized as defined in any of claims 13 to 15.
 - 21. A method according to claims 16 to 18, wherein said nucleic acids are labelled during or after amplification.
- 22. A polypeptide having an amino acid sequence encoded by a polynucleic acid according to any of claims 1 to 8, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in claims 2 or 3, and which contains at least one amino acid differing from any of the known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically

equivalent.

5

10

15

25

30

23. A polypeptide according to claim 22 comprising in its amino acid sequence at least one of the following amino acid residues:

115, C38, V44, A49, Q43, P49, Q55, A58, S60 or D60, E68 or V68, H70, A71 or Q71 or N71, D72, H81, H101, D106, S110, L130, I134, E135, L140, S148, T150 or E150, Q153, F155, D157, G160, E165, I169, F181, L186, T190, T192 or I192 or H192, I193, A195, S196, R197 or N197 or K197, Q199 or D199 or H199 or N199, F200 or T200, A208, I213, M216 or S216, N217 or S217 or G217 or K217, T218, I219, A222, Y223, I230, W231 or L231, S232 or H232 or A232, Q233, E235 or L235, F236 or T236, F237, L240 or M240, A242, N244, N249, I250 or K250 or R250, A252 or C252, A254, I255 or V255, D256 or M256, E257, E260 or K260, R261, V268, S272 or R272, I285, G290 or F290, A291, A293 or L293 or W293, T294 or A294, S295 or H295, K296 or E296, Y297 or M297, I299 or Y299, I300, S301, P316, S2646, A2648, G2649, A2650, V2652, Q2653, H2656 or L2656, D2657, F2659, K2663 or Q2663, A2667 or V2667, D2677, L2681, M2686 or Q2686 or E2686, A2692 or K2692, H2697, I2707, L2708 or Y2708, A2709, A2719 or M2719, F2727, T2728 or D2728, E2729, F2730 or Y2730, I2741, I2745, V2746 or E2746 or L2746 or K2746, A2748, S2749 or P2749, R2750, E2751, D2752 or N2752 or S2752 or T2752 or V2752 or I2752 or Q2752, S2753 or D2753 or G2753,

20 D2754, A2755, L2756 or Q2756, or R2757,

with said notation being composed of a letter representing the amino acid residue by its one-letter code, and a number representing the amino acid numbering as shown in Table 1,

or a part of said polypeptide which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one amino acid differing from known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.

24. A polypeptide according to claim 22 comprising in its amino acid sequence at least one of the sequences represented by SEQ ID NO 107 to 207 as listed in claim 5, or part of said polypeptide which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one amino acid differing from SUBSTITUTE SHEET (RULE 26)

5

10

15

25

known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.

- 25. A polypeptide having an amino acid sequence as represented in any of SEQ ID NO 1 to 106, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one amino acid differing from known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.
- 26. A recombinant polypeptide encoded by a polynucleic acid according to any of claims 1 to 8, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in claims 2 or 3, and which contains at least one amino acid differing from known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.
- 27. A method for production of a recombinant polypeptide of claim 26, comprising:
- transformation of an appropriate cellular host with a recombinant vector, in which a polynucleic acid or a part thereof according to any of claims 1 to 8 has been inserted under the control of the appropriate regulatory elements,
 - culturing said transformed cellular host under conditions enabling the expression of said insert, and,
 - harvesting said polypeptide.
- 28. A recombinant expression vector comprising a polynucleic acid or a part thereof according to any of claims 1 to 8 operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements.
 - 29. A host cell transformed with a recombinant vector according to claim 28.
 - 30. A method for detecting antibodies to HCV present in a biological sample, comprising:
 - (i) contacting the biological sample to be analysed for the presence of HCV with a polypeptide according to any of claims 22 to 26,

5

10

25

- (ii) detecting the immunological complex formed between said antibodies and said polypeptide.
- 31. A method for HCV typing, comprising:
- (i) contacting the biological sample to be analysed for the presence of HCV with a polypeptide according to any of claims 22 to 26,
- (ii) detecting the immunological complex formed between said antibodies and said polypeptide.
- 32. A diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one polypeptide according to any of claims 22 to 26, with said polypeptide being possibly bound to a solid support.
- 33. A diagnostic kit for HCV typing, said kit comprising at least one polypeptide according to any of claims 22 to 26, with said polypeptide being possibly bound to a solid support.
- 34. A diagnostic kit according to claims 32 to 33, said kit comprising a range of polypeptides which are attached to specific locations on a solid substrate.
 - 35. A diagnostic kit according to claims 32 to 34, wherein said solid support is a membrane strip and said polypeptides are coupled to the membrane in the form of parallel lines.
- 36. A pharmaceutical composition comprising at least one polypeptide according to any of claims 22 to 26 and a suitable excipient, diluent or carrier.
 - 37. A method of preventing HCV infection, comprising administering the pharmaceutical composition of claim 36 to a mammal in effective amount to stimulate the production of protective antibody or protective T-cell response.
 - 38. Use of a composition according to claim 36 in a method for preventing HCV infection as defined in claim 37.

5

10

15

20

25

- 39. A vaccine for immunizing a mammal against HCV infection, comprising at least one polypeptide according to claims 22 to 26, in a pharmaceutically acceptable carrier.
- 40. A vaccine according to claim 39, comprising at least one polypeptide according to claims 22 to 26, with said polypeptide being unique for at least one of the HCV subtypes as defined in claims 2 or 3.
- 41. A peptide corresponding to an amino acid sequence encoded by at least one of the HCV polynucleic acids according to any of claims 1 to 8, with said peptide comprising an epitope being unique to at least one of the HCV subtypes or types as defined in claims 2 or 3, and with said peptide containing at least one amino acid differing from any of the known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent.
- 42. A method for detecting antibodies to HCV present in a biological sample, comprising:
- (i) contacting the biological sample to be analysed for the presence of HCV with a peptide according to claim 41,
- (ii) detecting the immune complex formed between said antibodies and said peptide.
- 43. A method for HCV typing, comprising:
- (i) contacting the biological sample to be analysed for the presence of HCV with a peptide according to claim 41,
- (ii) detecting the immune complex formed between said antibodies and said peptide.
- 44. A. diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one peptide according to claim 41, with said peptide being possibly bound to a solid support.
- 45. A diagnostic kit for HCV typing, said kit comprising at least one peptide according to any of claim 41, with said peptide being possibly bound to a solid support.

SUBSTITUTE SHEET (RULE 26)

- 46. A diagnostic kit according to claims 44 or 45, wherein said peptides are selected from the following list:
- at least one NS4 peptide,
- at least one NS4 peptide and at least one Core peptide,
- at least one NS4 peptide and at least one Core peptide and at least one E1 peptide, or,
 - at least one NS4 peptide and at least one E1 peptide.
 - 47. A Diagnostic kit according to claims 44 to 46, said kit comprising a range of peptides which are attached to specific locations on a solid substrate.
- 48. A diagnostic kit according to claims 44 to 47, wherein said solid support is a membrane strip and said peptides are coupled to the membrane in the form of parallel lines.
 - 49. A pharmaceutical composition comprising at least one peptide according to claim 41 and suitable excipient, diluent or carrier.
- 15 50. A method of preventing HCV infection, comprising administering the pharmaceutical composition of claim 49 to a mammal in effective amount to stimulate the production of protective antibody or protective T-cell response.
 - 51. Use of a composition according to claim 49 in a method for preventing HCV infection as defined in claim 50.
- 52. A vaccine for immunizing a mammal against HCV infection, comprising at least one peptide according to claim 41, in a pharmaceutically acceptable carrier.
 - 53. A vaccine according to claim 52, comprising at least one peptide according to claim 41, with said peptide being unique for at least one of the subtypes or types as defined in claims 2 or 3.
- 25 54. An antibody raised upon immunization with at least one polypeptide or peptide SUBSTITUTE SHEET (RULE 26)

according to any of claims 22 to 26 or 41, with said antibody being specifically reactive with any of said polypeptides or peptides, and with said antibody being preferably a monoclonal antibody.

- 55. A method for detecting HCV antigens present in a biological sample, comprising:
- 5 (i) contacting said biological sample with an antibody according to claim 54,
 - (ii) detecting the immune complexes formed between said HCV antigens and said antibody.
 - 56. A method for HCV typing, comprising:
 - (i) contacting said biological sample with an antibody according to claim 54,
- 10 (ii) detecting the immune complexes formed between said HCV antigens and said antibody.
 - 57. A diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one antibody according to claim 54, with said antibody being possibly bound to a solid support.
- 15 58. A diagnostic kit for HCV typing, said kit comprising at least one antibody according to claim 54, with said antibody being possibly bound to a solid support.
 - 59. A diagnostic kit according to claims 57 to 58, said kit comprising a range of antibodies which are attached to specific locations on a solid substrate.
- 60. A pharmaceutical composition comprising at least one antibody according to claim 54 and a suitable excipient, diluent or carrier.
 - 61. A method of preventing or treating HCV infection, comprising administering the pharmaceutical composition of claim 62 to a mammal in effective amount.
 - 62. Use of a composition according to claim 60 in a method for preventing or treating HCV infection as defined in claim 61.

Figure 1		1/74	50
HCV-1	1a .	ATGAGCACGAATCCTAAACCTCAAAAAAAAAAAACAAACGTAACACCA	ACCG
HCV-J	1b	A	
HCG9	1c	G	
BNL1	1d	C	
BNL2	1d	C	
CAM1078	1e	GGA-A	
FR2	1f	CC	
76	2-	AAGCA-A	
HC-J6	2a	AAA-AA-AAA	
HC-J8	2D	AA	
S83	2c	AT	
NE92	2d	AT	
FR4	2f	AT	
BNL4	2e		
BNL5	2h	AT	
N77 1	25	ACTGCA-A	т
NZL1	24 25	ACTACTA-AACT	r
HCV-TR	30	ACTACGCA-A	T
NE48	3C	ACTACTACGCA-A	T — — —
NE274	3a -	ACTACGCA-A	T
NE145	3e	ACTAGCA-AG	r
NE125	3f	ATTACC	J
2.4	4a -	CC	
21	4h	AGC	
GB358	40	GC	
	40.	C	
DK13	40		
GB809	4e	GC	
BNL7	4 K		
BE95	5a -		
HK2	6a -	ACTACGCA-A	· ·
FR1	75.	ACTACGCA-ATT	
EKI			
VN4	8a -	ACTAT	ſ
VN13	8b ·	ACTGCA	
11120			
VN12	9a ·	ACTACGCA-AA	
NE98	10a -	ACTAGCA-A	N

Figure 1 -	-continued		2/74			
	51					100
HCV-1	la TCGCCC	CACAGGAC	STCAAGTTC	CCGGGTGGC	CGGTCAGA	TCGTTGGTGGAG
HCV-J						
HC-G9						C
BNL1	1d C	TK-GS-	 			
BNL2						
CAM1078						
FR2						G
	•			_		
HC-J6	2a	A	-TT-	C	C	C
HC-J8						C
S83						C
NE92						
FR4						C
BNL3						C
BNL5	2h C		-T	·СТ	C	Č
						_
NZL1	3a				A	
HCV-TR						
NE48						
NE274						
NE145						
NE125						
Z4	4a C	CAT	-A	T	C	C
Z1	4b	CATT-	-GA	C	C	C
GB358	4c C	CAT	-T	CT	c	C
DK13	4d C	-AT	-T		C	C
GB809	4e C	CAT	-T	T	C	C
BNL7						C
BE95	5a			CT		C
HK2	6a	-AC				C
FR1	7a	TAT		C	C	
VN4	8a C				C	
VN13	8b					
	_					
VN12	9a	-ATT-		C		
00						
NE98	IUa CG		-T	-AC		

Figure 1 -		
HCV-1 HCV-J HC-G9 BNL1 BNL2 CAM1078 FR2	101 1a TTTACTTGTTGCCGCGCAGGGGCCCTAGATTGGGTGTGCGCGCGACGAGA 1bC	O
HC-J6 HC-J8 S83 NE92 FR4 BNL3 BNL5 NZL1 HCV-TR	2a -A	
NE48 NE274 NE145 NE125	3c -AGTAC-T 3d -CACAGTTC-T 3e -AACACAGTTC-T 3f -AG-AACAGT-C-T	
Z4 Z1 GB358 DK13 GB809 BNL7	4a	
BE95 HK2	5aTC-G 6aCC-G	
FR1	7aC-T	
VN4 VN13	8a -CGC-C	
VN12	9a -C	
NE98	10aGC-AA	

Figure 1 -	continued	4/74			200
HCV-1 HCV-J HC-G9	1a AAGACTTCCGAGG		TAG	AA	TATCCCCAA
BNL1	ld	A	TCG	A	
BNL2	1d				
CAM1078	leG				
FR2	1f				
			C A G	- A	
HC-J6	2aG		TNC		· m
HC-J8	2bT				
S83	20	-ACGG-	-1AC		,G
	2cAA-	CGA-	- <u>-</u>	CC	:T
NE92	2dA	CGA-	·-TGG	CC	
FR4	2fTA-	CGA-	·-TAG·	CC	A
BNL3	2eTA-	CGA-	-TAG-	CC	T
BNL5	2hAA-	CGA-	-TGG-	CC	T
NZL1	3aATA-	AG	-CAC	A	
HCV-TR	3b				
NE48	3c				
NE274	3dA	NGC-	-C770CC-C-	G	
NE145	3eA	AGC-	-CAACC-G-	G	
NE145 NE125	3fAT	AC-	-CAC-G-	A	1
NEIZS	21WI		-CAC-G-	G	
17 A	4.5			_	
Z4	4a		-TCG-	A	
Z1	4bG				
GB358	4cG				
DK13	4dG				
GB809	4eG				
BNL7	4kG		-TG	-CA	
BE95	5aGA	C	-TAC-G-		T
		_			_
HK2	6a	-ACGCA-	CG-	-CA	AA
			• •	•	••
FR1	7aCA	CGA			22
• • • •	,	O n	0		AA
VN4	8aTA	CCC	_	C 7 7	*
·					
VN13	8bATA	CGCA-	-6	-CA	AG
					_
VN12	9aGA	CGG-CA-	G-	-CAA-	A
NE98	10a		-CAG-	-CAC-	G

Figure 1 - continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 CAM1078 FR2	201 250 la GGCTCGTCGGCCCGAGGGCAGGACCTGGGCTCAGCCCGGGTACCCTTGGC lbT	כ
HC-J6 HC-J8 S83 NE92 FR4 BNL3 BNL5	2aAGCTACTAATGAA-AAAC 2b A-AGCTACCA-TGAAAAT 2c A-AGCAACTA-TGAAGAA	
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a GAGACT	
Z4 Z1 GB358 DK13 GB809 BNL7	4a GC-AA	
BE95 HK2	5aGC-AACCTGAA	
FR1	7aTAC-AGACAC-T-GGAC	
VN4 VN13	8a A-TGC-AC-AAACC-TCC 8bTGAC-AAACC-TA	
VN12 ·	9aTGC-A-AA-C-AC-ATT	
NE98	10aGCAATT	

Figure 1- continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 FR2	251 la CCCTCTATGGCAATGAGGGCTGCGGGTGGGCGGGATGGCTCCT lb	A C
HC-J6 HC-J8 S83 NE92 FR4 BNL3 BNL5	2aA-C-GACTCA	C
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a	C C C
Z4 Z1 GB358 DK13 GB809 BNL7	4a	C AT CT
BE95 HK2	5aTC-CCTAGGC 6a -TTACTAT	
FR1	7aTCA	
VN4 VN13	8a -TTATTAC 8b -TTGTTCAG	C
VN12	9aTGC	
NE98	10aAG	C G

Figure 1 - continued

HCV-1	301 1a CGTGGCTCTCGGCCTAGCTGGGGCCCCACAGACCCCCGG	350 .
HCV-J HC-G9 BNL1 BNL2	1bT	GA
FR2	1fCATA	AA
HC-J6 HC-J8	2aATCTCTCTAT 2bCGTCT	-AC -AAA
\$83	2cCTCTCA	-AAA
NE92	2dAGCGTCAAT	-ACA
FR4	2fGCCTCGA-AC2eA	-ACA
BNL3 BNL5	2hA	
NZL1	3aCATCA-AT	GC
HCV-TR	3bA-AT	AC
NE48	3cCTGA-AT	AAC
NE274	3dCATCTAT	AT
NE145	3eCCAGTAC	AC
NE125	3fA-AT	
Z 4	4aCATCTA-ATT	GA
Z1	4bCTCAGTCTATT	C
GB358	4cA-GTCTA-ATT	AC
DK13	4dGTCTG-ATT	GC
GB809	4eCGGTCTT-ATT	G
BNL7	4kCT	
BE95	5aAATAT	
нк2	6aCACATAT	
FR1	7aCGTATAC	
VN4	8aCA-ATA-AC	GC
VN13	8b -NCCATT-AT1	.1-GC
VN12	9aCGGANAT	И-GC
NE98	10aC	

Figure 1 - continued

HCV-1 HCV-J HC-G9 FR2	351 400 1a CAATTTGGGTAAGGTCATCGATACCCTTACGTGCGGCTTCGCCGACCTCA 1b T
HC-J6 HC-J8 S83 NE92 FR4 BNL3	2aCG
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a
Z4 Z1 GB358 DK13 GB809	4a C
BE95	5a TT
HK2	6a GTTTTT
FR1	7aCA-NNC-A
VN4 VN13	8aCACT
VN12	9aCC

Figure 1 - continued

HCV-1	401 45 1a TGGGGTACATACCGCTCGTCGGCGCCCCTCTTGGAGGCGCTGCCAGGGCC	:
HCV-J	1bTT	
HC-G9	1cAGAT	ı
FR2	1fTAA	
HC-J6	2a	
HC-J8	2bTCAT	,
S83	2cCGTTCGCA	
NE92	2dTGAGT-TCAT	
FR4	2fTGG-G-CTA	
BNL3	2eNCG-TGG-GCG-TN	
NZL1	3a	
HCV-TR	3bG-G-GTCAA	
NE48	3cT	
NE274	3dTTTTTTTTT	
NE145	3eT-TTGG-ATCG	
NE125	36TCG	
145123	3fTCA	
Z 4	4aACAG	
21	4bATAG-GTTC	
GB358	4cACG-GTTC	
DK13	4dACGA	
GB809	4eA	
BE95	5aTC	
HK2	6aTCGGT-GCTCGGCTG	
FR1	7a	
VN4	8aTCTGATGW-GTCGGN	
VN13	8b -A-AT	
*****	OD - R R1	
VN12	9aATGTCTGGCAA	

WO 96/13590

10/74

Figure 1 - continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 FR2	451 500 1a CTGGCGCATGGCGTCCGGGTTCTGGAAGACGGCGTGAACTATGCAACAGG 1bATTA-ACTC 1dN-ATCN-GTNNNNNNNNNNNNNNNN
HC-J6 HC-J8 S83 NE92 BNL3 FR4 BNL4 BNL5 BNL6	2aC
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a CGA
Z4 Z1 GB358 DK13 GB809 BNL7 BNL8 BNL9 BNL10 BNL11 BNL11	4a
BE95	5aCACTGACTGGA
HK2	6aCAGACAA-CGGA-CT
FR1	7aTACAA-CGGCTC
VN4	8a TN-AN-NCA-CGN-ATCN
VN12	9aNATACCA-CGGA-A
NE98	10aAA-TT-TC

Figure 1 -continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 FR2	501 550 la GAACCTTCCTGGTTGCTCTTTCTCTATCTTCCTTCTGGCCCTGCTCTCTT lbT-G-CT-GC-T-T-AC- lcC-C-CT-GC-TTGC- ldT-GC	
HC-J6 HC-J8 S83 NE92 BNL3 FR4 BNL4 BNL5 BNL6	2a T-AC	
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a T-GC	
Z4 Z1 GB358 DK13 GB809 BNL7 BNL8 BNL9 BNL10 BNL11	4a TC	
BE95	5aTT-ACG-	
HK2	6aTCC	
FR1	7aT	
VN4	8aTG-	
VN12	9aT	
NE98	10aTT-AA-	

Figure 1 - continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 FR2	551 1a GCTTGACTGTGCCCGCTTCGGCCTACCAAGTGCGCAACTCCACGGGGCTT 1b -TCA-CAC-TG-GGTGT-CA-A 1cCACTGT-GGTTG-G 1dG-TAA-KA-CTCG-GG-AT-CG-A 1fC-CACA-CTTG-GA-G-A-AC-ATGGC
HC-J6 HC-J8 S83 NE92 BNL3 FR4 BNL4 BNL5 BNL6	2aA-CCACCG-TCCTGC-GAAGATGTACCGGC 2bG-CAA-TGTAGTGGCA-GATT-GTTCTAGC 2cA-CTA-TCGTGG-GCAAGGAGGC-ACTCC 2d -TA-CG-TCC-GTGGCAAGAGCA-CTC- 2e -TG-CCT-TCT-N-GTTG-GCAAATAGTCA-GCC 2f -TA-CCTGTATAGTAAGAAGCCACT-C 2g -TG-CCT-TCTGTGGTAAGAGTACCA-G 2h -TC-CG-GCTGTGGCAAGAGCCACTC- 2iA-CCG-TCTGTGTGCGCGGTTTC-
NZL1 HCV-TR NE48 NE274 NE145 NE125	3aA-T-CATAAG-CAGTCTAG-GTGGTA-GT-TCC 3bTGCGT-GTAG-GTACACGA-GT-TCA 3cGTCTGTTAG-A-GGCT-G-GTACGTGTAT-CCC 3dGTCTGTTG-A-GGATTGTACGTGTGT-TCC 3eCT-TGCTAGTC-GG-TGG-GTG-AT-CTC 3fGT-TCCAGGGCTAG-GTACA-GA-GT-CCA
Z4 Z1 GB358 DK13 GB809 BNL7 BNL8 BNL9 BNL10 BNL11	4a
BE95	5a -TCCTGCTAGTT-CCTACATGT-TA
HK2	6aC-CAACATCTTACCTACGGTA
FR1	7aC-CACAACAAATTCAAGGT-TA-C
VN4	8aC-TAACAACCGGCGTTATACAAGT-TCG
VN12	9aC-CCACTCCACTAA-CTATGCTAAGT-TG
NE98	10aCT-ACAA-AG-C-GGCTGG-GTACTTGT-CAC

Figure 1 - continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 FR2	601 1a TACCACGTCACCAATGATTGCCCTAACTCGAGTATTGTGTACGAGGCGGC 1bTGCCT-CATA 1cTCTT-CCCA-CTATA 1dTTCTT-CCCA-CTATA 1dTTCCTT-CCCA-CTAT-AG 1fTTCTT-CGGCCCA-TAAA
HC-J6 HC-J8 S83 NE92 BNL3 FR4 BNL4 BNL5 BNL6	2a ATGGCCA-CTGATCACC-GGC-ACTCCA 2b TCTT-AAACCCACC-GGCCTCA- 2c ATGCCGCT-CT-CT-GGCCTT-A 2d ATG-ACAGAGTCCC-GGCCTCAG 2e -TATG-CACT-CAACCCA-GGC-ATTN 2f ATG-CGTC-TG-C-TGACCCC-GGCCTCAG 2g ATG-CAC-TT-CAACCCA-C-GGC-AAT-CA 2h -TATG-GT-AAGCCC-GGCCTTAA 2i ATG-GT-GAGCCC-T-GGCCTC-A
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a GT-C-TCCTT-CTAGCTC-A 3b TGTGC-TCCTTGGC
Z4 Z1 GB358 DK13 GB809 BNL7 BNL8 BNL9 BNL10 BNL11 BNL11	4a -TA
BE95	5aTTT
нк2	6aTCACCCCTGA
FR1	7aTC-TCT-GAACCC-T-TTA
VN4	8aTCAGCCCTTA
VN12	9aTTC-ACTAGCCTAGC
NE98	10aATGATCCAGGGTCTC-G

VN12

NE 98

```
Figure 1 - continued
                             14/74
               651
                                                         700
  HCV-1
            1a CGATGCCATCCTGCACACTCCGGGGTGCGTCCCTTGCGTTCGTGAGGGCA
  HCV-J
            1c GA-CCTG---A----TCTG--C----T--G--C-A---A--C-----
  HC-G9
 BNL1
           1d --G-ATG---A----TAC--A-----G--C---AT-
 BNL2
           1d T-G-ATG---T----G-C--A----T--G--C-----G---AA--
           lf G--CAT-----T----G--T----N--G--C---A-A--G--A---
 FR2
 HC-J6
           2a G-C---TG----C---GTC--C-----G----AGAAA-T---G-
           2b T--C--AG-T--C--TCT---T-A----A--T-AGAA---TAATG
 HC-J8
 S83
           2c A-GA--AG-G--T--T--T--A-----T-AG---ACC-C--
           2d G----TG-T--T---GTC--C----T-AGGAGA-----
 NE92
 BNL3
           2e G--C--GG-G--T--TGT---T-A--T----C---AGAA-AGCTC-G
           2f G--C--GG-G--C--TGT---T--A--T----C--T-AGA-GTCA--T-
 FR4
 BNL4
           2g G-GC--GG-G--T--TGT---T-A--T----G--T-AGTTGC-----
           2h G----TG-G-T---GTC--T-A-T-T-A-T-AGA-GC-CCAA-
 BNL5
 BNL6
           2i G--G---G---T--GTC--T--A--T--T--C--T-AGT-GA---A--
           NZL1
           3b A----TG---TTA--C--A----G--C----CACAACC----
 HCV-TR
           3c -C---T----TTG--C--T----A--C-----C-AAA-CAAT-
 NE48
           3d T--A-T----TTG--A--T--T--G--C-----AATCA----
 NE274
           3e A----TG------TG--T--T----T--C-----G-AGA-C----
 NE145
           3f TA---T------TG--C--C--T--G--C---AC---C---T-
 NE125
24
          4a -C--CA----A---TTG-----A--C--T--GATGACT--G-
\mathbf{Z}\mathbf{1}
          4b GC-CCA---A----TTG--A----T----C--T--G--GAC--AG-
GB358
          4c GC-CCA----A---CTC--A----TT-A--C----GA-G-TT--G-
          4d TT-CCA----T-A---CTC----A----T-----GA-G--A--G-
DK13
          4e -A--CA----T-A---CTC--A-----A--C--T--GAAGACC--G-
GB809
BNL7
          4k -C--CA----T-----G--C-----GA-A------G-
BNL8
          4k -C-CCA----T-----CT---A--T-----GA-CT--G-
          4k -C--CA----T----TCTC--A--T----G--C-----GA-A-T---G-
BNL9
          4k -C--CA---T-AGCACT---A--T----G--C----GA-A-T---G-
BNL10
BNL11
          4k -C--CA----T-----CT---A--A-----G--C-----GAAA-----A-
          41 -C--CA----T-A---CTA--A----T-A--C--T--GAAGACT--G-
BNL12
BE 95
          5a TA-CCTG----A---G-A--T--T----G----T--CATGACA--T-
HK2
          6a T-C-ATG---T---TTTG--T--A---T-G---T--GA-G-TC-ATG
         7a GACCATG--A----TCT---A--T--TA-CAAG-C---G-
FR1
         8a GACACTG--TT-----TTG--T----T--GAAGRT-RA--
VN4
```

9a T-GCATG-----TCTC----T----C----GAAGACC----

10a G---ATT-----C--TTA--T--C--T-----A--CTCT----

Figure 1 continued

```
701
                                                            750
           1a ACGCCTCGAGGTGTTGGGTGGCGATGACCCCTACGGTGGCCACCAGGGAT
 HCV-1
 HCV-J
           1b -TTT---CC-T--C---A---C-C--T--C--C--GG-----A-C
           1c -----CT-CC-T-GT--C--C-A---G-----
 HC-G9
 BNL1
           1d --CATCTCC-C--C--A----C-C---C-T--GGT--AAA-Y
           1d --T-T--TC-T--C--A--C-RC-C----C-T--GGT--AA--C
 BNL2
 FR2
           1f -TAT---CC-T--C----AC--C-C----C--C--AG-GC--ATC
           2a -TA-A--TC----C--A-AC--G-CT-A--G-AT-----GTGCA-C-G
 HC-J6
 HC-J8
           2b G-A---T-CAT--C--A-ACAAG-A--A-C-AC---TGTG-AAC-C
 S83
           2c ---T---TC-A-------C--G-TG----C-ATC-C----TA--TC-A
NE92
           2d --ATA--CC-C----A-AC--G-TT-G--C-ATA-A--TGTG--CC-A
 BNL3
           2e GTCGG-TCCAC----A-CC----CT-G--C-ACA-A---GTG--CA-A
 FR4
           2f -TAGGA-CTTC-----ACA---G-CT-G--C-AC----TGTG--CCGA
BNL4
           2q -TAAG--CC----C--A-AC--G-C--T--C-AC----TGTG-ACC-G
BNL5
           2h -TCAG--TC-C--C--A-AC-TG----A--C-AT-----GTG--CC-A
           2i --A----CC-C--C--A-AC--G-C-----ACA-C--TGTG--CC-A
BNL6
NZL1
           3a -TA-A--T-C---C---ACCC-AG----A----AGT----T-C
HCV-TR
           3b -- CAA--ATCA--C---ACAA--G-CT-AA-G------GTT---ACC
NE48
           3c --A--A---C---C---A-AC--G----T--G--A-----GGT---TC-C
NE274
           3d --T----CAA--C---A-TC--G--G-A--A--A--A-----GGTT-A-T-C
NE145
           3e --A-A---GA---C---ACCC--GC---A--A--A-----AGT---AT-C
           3f -- CAG--A----C--AC-C-AG-A--A--G--A---TGT--AAC--
NE125
24
           4a --A-A---C-T--C---AC-C--G----G----A----TGT-GCAC-C
21
           4b -TA-T--TC-C--C------C-CT----------G-GCCCT--
GB358
           4c -TCAG--AC-C--C-----CC-C--T--C--C-----GG-GCCTT-C
DK13
           4d --AAG--T-CA--C----T-TC-C-----TG-GCAAC--
GB809
           4e --CAG---C------CC-C--T--C-A-----GT-GCCTT-C
BNL7
           4k -TCAG--AC-T--C----A--CC-T-----C--C--AG-GCCAT-C
BNL8
           4k -TCAG--AC-T--C-----CC-T--T-----C--AG-GCCAT-C
BNL9
          4k -TCAG----T--C-----CC-T-----CA-C--AG-GCCAT-C
BNL10
          4k -- CAG--AC-C--C------CC-T-----C--C--AG-GCCAT-C
BNL11
          4k -TCAT--AC-C--C------CC-T-----C--C--AG-GCCAT-C
BNL12
          41 --A-T---C-C--C------CT-A--A-----C-----G-GCCCATA
          5a -T-TGAGT--A--C----CCAA--T-----AC--T-AG--CC-AGC
BE95
HK2
          6a -TCGG--C-CC-----TACCAA--
          7a -T-AG--AC-A-----C-CC-TG-CT----C-CT-A---GT-CCCA-C
FR1
VN4
          8a -TCAA--CC----CA-GCCT----G--CC---AGTGCC-A-C
          9a --CTGA-C-A-----C--T--GCCT----G--AT----GGTGCA-A--
VN12
         10a -TA-A--A--C--A-CC-TG---G--Y--C--GTG-A-TCG
NE98
```

Figure 1 - continued

```
800
             1a GGCAAACTCCCCGCGACGCAGCTTCGACGTCACATCGATCTGCTTGTCGG
  HCV-1
  HCV-J
             1b A---GCA----A-C---ACAA-A-----C---G-----T----C--T--
  HC-G9
             1c TCGCGCG-----TC-GTG--G----GTG----CTC-A----
  BNL1
             1d -CT-GTG----A-TR--GCAA-C-----G---CT-----T--
             1d -CT--TG----TA-TG--GCAA-C----CT-TG----CT----G--T--
  BNL2
  FR2
             1f -CG--CGCT---ATCGATG--G-G--G--C--C--C--C--G--
            2a CC-GGCGC--T-A--CA-GGCT-A--GACG----T--CA--G----GAT
 HC-J6
 HC-J8
            2b C--GGTGCG-T-A-TCGTAGC--G---ACA---G---CA--A-C--AAT
            2c CCTGGCGCT-T-A-T-A-GGC--G---GCA-----A-CA-C--GAT
 S83
 NE 92
            2d CCTGGTGCG-TTA-C-A-GGC--G--GACG--T--T---ACCA-CA-T-C
 BNL3
            2e CCTGGTGCT-T-A-C-A-GGA--G--GGCA-G--T---GCCG-C--GAT
            2f CCTGGTGCT-T-A-T-GAGGT--G--GGC----T---ACCA-C--GAT
 FR4
 BNL4
            2g CC-GGCGC--T-A-T-G-GGCT-G--GACG----T--CACCA-C--GAT
            2h CCTGGCGCG-T-A-C-G-GGTT-G--GACG----T-CACCA-C--T-C
 BNL5
            2i CCTGGCGCG-TTA-C-A-GGC--G--GACA--T--T--CA-CA-----C
 BNL6
 NZL1
            3a -T-GG-GCAA-TA-TG-TTC-A-A--CA----TG-G--C--AT-A--A--
            3b CTTGGCG-GA--A-CG--TC-A-C--ACC--TG-G---A---G--A--
 HCV-TR
 NE48
            3c -T-GGTGCGA--A-CG-ATC-A-C--CG-G---G-G-----G-G--
 NE274
            3d -CTGGCGCGA--A-TG-ATC-A-C--CA----TG-G-------G--G--
 NE145
            3e -CTGGTGCAA-GA--G-TTCCG-A--CG-A---G-G---T----A----
 NE125
            3f CCTGGCGCAGT-A-CG-ATCAA-C--CA-G--TG-G---T--A-G--G--
            4a CCGGGCGCT--GCTTGA-TC-T-C--G--A--TG-G--CT-AA-G--A--
 24
            4b CC---CGCA--GTTAGA-TCCA-G--CA-G--TG-A--C--A-G--G--
 21
 GB358
            4c AT-GGCGCT--GCTTGAATCC--C--GA----TG-G-----A-G--A--
 DK13
           4d CTG--TGCT--GCTTGA-TCTT-GA-----G-G----A-G--G--
GB809
           4e -T-GGTGCT--GCTCGA--CCT-G--G--C--TG-G--C--A-G--A--
BNL7
           4k AT-GGCGCG--ACTTGA-TCT--A--GA----TG-G--CT--A-G--G--
BNL8
           4k AT-GGCGCA--GCTTGA-TCT--G--GA----TG-G-----A-G--G--
BNL9
           4k AT-GGCGCA--GCTTGA-TCCT-G--GA----TG-G----A-G--G-
BNL10
           4k AC-GCGGCG--GCTTGA-TCC--G--GA----TG-G-----A-G--G-
BNL11
           4k AT-GGCGCG--ACTTGA-TCT--A--GA----TG-G---G--A-G--G--
BNL12
           41 CTTTCGGCT--ACTT-T-TCCG-A--G--G--TG-G-----A-G--G--
BE95
           5a CT-GG-GCAGT-A--G-T-CT----GA-AGC-G-T--CTAC--A-CG--
HK2
           6a -CTTCCACG----A---GGAT-C--CA-G--TG-G----T----CG--
           7a TCATC-G-G--AATCCACGG-T----C-A---G-A--C--C--T--
FR1
           8a -CGTCTACG--A-TC--CGG-T-C--CAAA--TG-G--CA-CA-G--G--
VN4
           9a -CGTCGG-GT--ATC-G-GGTG-C--CGAG---G-G--CT-G--G-
VN12
NE98
         10a CC-TGCGC-G--A-CG-CTCT--C--CACG---G-G--A--A-G--G--
```

```
Figure 1 - continued
```

HCV-1 HCV-J	801 850 1a GAGCGCCACCCTCTGTTCGGCCCTCTACGTGGGGGACCTATGCGGGTCTG 1b -GCGTG-TCTA-GTCAC-
HC-G9	1c -GCTG-GTTA-GT-ACCA
BNL1	1d -G-NNGTCTA-GRT
BNL2	1dCAG-GT-TCCTA-GCAC-
FR2	1f -GCAGTGTCAA-GA-TTTGGC-
HC-J6 HC-J8	2a -TCGCCTT
583	2b -GCATGGCCT-GTATG-GG-C-
NE 92	2c -TCTTGGTTTG-GTCG-GC
BNL3	2d ATCTGT-TCTGA-A-AAGTCG-G-
	2e -TCCTGATG-GCG-A-
FR4	2f -TC
BNL4	2g -GTTGTAA-CG-GTCG-G-
BNL5	2h -TCTTGCATT-GTCT-CCG-A-
BNL6	2i -TCGTCTT-GT
NZL1	3a CGCGGA-GCTGTTA-GTG
HCV-TR	3b CGCACGACAAGGGCGCT-TG
NE48	3c T-CGTAT-GATCTTC-A-
NE274	3d AGCTTGT-GCCGGTTCTA-GTAG-C-
NE145	3e CTTGCGTCTTC
NE125	3f TGCAGGAATT-ATT-GG
Z4	4a CGCGTT-GTTTTCAGG
Z1	4b TGCGTTA-GCTA-TA-TGTAGGC-
GB358	4c TGCTTGCGCCTTA-CAGTGGC-
DK13	4d CGTCCA-CAG-GTGG
GB809	4e TGCTG-GCCCCTGGCT
BNL7	4k -GCTG-TATA-CTT-RTYGGCT
BNL8	4k -GCTTG-TC-ATA-CTT-GTCGGCT
BNL9	4k -GCGTGATA-CTT-GTCGG
BNL10	4k AGCTTG-TATA-CYT-GTCGCT
BNL11	4 k -GCTTG-TAT-G-CT
BNL12	41 TGCATA-CGTT-ACGG
BE95	5a AG-GTGCCGT-AAGCGTG-AC
HK2	6a CGCAGTGG-TCATGA-CGTCC
FR1	7a -GCAGG-AT-TA-GA-CA-CTTAGCA
VN4	8a CGCTG-GTATA-GTGGCC
VN12	9a TGCTTG-GTCTA-GCTTGGGC
NE98	10a RGCGACATAAATAG-GC

Figure 1 - continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 FR2	851 900 1a TCTTTCTTGTCGGCCAACTGTTCACCTTCTCTCCCAGGCGCCACTGGACG 1b -TCTCGATC-CGT-TGA 1cCTGA-CATC-C-ATGCATA 1dCC-CTG-AT-AC-CATGCATA 1dCC-TG-AT-AC-CTTGTCATA 1fCC-TGTAGTCG-T
HC-J6 HC-J8 S83 NE92 BNL3 FR4 BNL4 BNL5	2a -GA-GCA-CGATTGGACAATTT 2b -GA-GAC-ATCGGGCTTGG-AAACAAAACTTC 2c -GA-GG-CCTGG-CGGT-G-GGACAA-ATAC-TTT 2d -GA-GT-G-CTTCTG-CT-AGCAATTAA-TTT 2e -GA-GA-A-CT-CAGGCTT-G-GG-AG-AT-ACTTC 2f -GA-GA-A-CA-CGG-TGC-GT-GAGCAATATACTTTT 2g -GA-GA-A-CT-CTGG-TGTTGGGCAA-ATAACTTT 2h -GA-GT-GTCTT-TTGACTCAAATCTTC
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a C-G-AGCCGAGATC-ATCAA 3b -GG-AGCAGATC-CACC 3c -TCC-A-AGCAAAGAC-ACAA 3d CT-G-G-AGGCTAGATC-T-AGAAC 3e CGG-GCCT-AAGGTC-T-T-TACT 3f -TCGGCTAGAG-TCAAT-ATC
Z4 Z1 GB358 DK13 GB809 BNL7 BNL8 BNL9 BNL10 BNL11 BNL11	4a CCGA-GGGA-ATTCGGGC-TC 4b CAGGGACGAGC-CGC 4c -AT-G-T-T-GAT-TCAGGCT 4d -GCT-GTCAATC-CC 4e -ACT-G-AACAAGC-AT 4k -GCT-G-T-T-GATTCGA-AT 4k -GCT-G-T-T-GAT-TCGA-AC
BE95	5a -ACT-GAAATAGGTC-C-AGGCT
HK2	6aT-G-CGATCAGC-C-TTT
FR1	7a -AA-CT-GAGGTTTAGGT-A-TATCA-GTT
VN4 VN12	8a -TCCTAGCGCAGGTCATGTCA-GTT
NE98	9aATGT-TGATC 10a -AYGGGGT-A-GGAGA-ATC-C-AGTT

Figure 1 - continued

```
950
 HCV-1
          1a ACGCAAGGTTGCAATTGCTCTATCTATCCCGGCCATATAACGGGTCACCG
 HCV-J
          HC-G9
          1c -----G-G--A----T--
          ld ----A---
 BNL1
 BNL2
          1d --A--G-AG-----C----A---
          FR2
 HC-J6
          HC-J8
          2b --C---AG-----C--T--C-AA--T--C--C--C--T--
          2c GTC--G-AA-----C--T--C--A--C--G----GC--T-----A----
 S83
 NE92
          2d GTC--G-AC-----C--T--C--A---C--A----C--C--T--A--T--
 BNL3
          FR4
          2f GTC--G-AA-----C--A--C--A------C--A--A--T--
BNL4
          2q T-C--G-A-----T--C---
BNL5
          2h GTC--G-A----G--A
NZL1
          3a GTC--GACC--T--C-----GC-G--C--A-----C-TT-A--A--T--
HCV-TR
          3b GT---GACG-----C----G--A--C--A-----G-TT-A--A--T--
         3c GTT--GCA-----C----AC-G--C--A--T---G-TT-A----T--
NE48
NE274
         3d GT---GACC------AC-G--C--T--T--C---T-A--A---A-
NE145
         3e GTC--GACC-----C----GT-G--C--A--------C--A--A--T--
NE125
         3f GTC--GTTG-----AC-A--C--A--A--C--T--A--A--T-A
Z4
         4a ----G-AG------T--C-----CA-T-----C--C--A-
Z1
         4b --C--G-A-----C----C----T--T--CG-CT----C--A-
GB358
         4c ----G-AC-----T--C----CG-G--G--CG-T-----C--A-
DK13
         4d --C----AC------T--C-----CA-A--A--A---C--A--A--A-
GB809
         4e --C--G-AC--T----T--C----CG-A--G----T----C--T--
BNL7
         4k --T----A-----T--C---
         4k G-C--G-A-----T-----
BNL8
BNL9
         4k --C----A-----C----
BNL10
         4k --C--G-A-----T--C---
         4k --C--G-AA-----T--C---
BNL11
BNL12
         41 GTC----AC-----C--T--C---
BE95
         5a GT---GAAC-----C--T--CAGT------G-T--C--C----
HK2
         6a GT----AC----C----C----A-A----CG-C--C--C--A-
FR1
         7a --C--G-A---T--C-----NA-CN-T-----CG-C----A---A-
VN4
         8a GTC--G-AG--T--C--T--CA-A--G----C--T--A----
VN12
         9a G-C--G-AC-----C--T--C------G-A-----C--C--T--G-----
NE98
        10a GTC--G-AC----C--T--C---
```

Figure 1 -continued

HCV-1 HCV-J HC-G9 FR2	951 957 la CATGGCA lbT lc AT lf NNNNNN
HC-J6 HC-J8 S83 NE92 BNL3 FR4	2aG 2bT 2cG 2d GG 2eG 2f ANN
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a AT 3b TG 3c GT 3d GT 3eT
Z4 Z1 GB358 DK13 GB809	4a GG 4b GC 4c G 4d AT 4e GT
BE95	5a G
нк2	ба GТ
FR1	7a G
VN4	8a A
VN12	9a G G

_						_
F	3	$\boldsymbol{\sigma}$	11	*	0	ッ
•	_	•	•	_	-	_

HCV1 HCV-J	la 1b	1 50 MSTNPKPQKKNKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATR
BNL1	1d	R-TXXXXXX
BNL2	īd	R-TXXX
CAM1078	1e	VA-
FR2	1f	R-T
нсј6	2a	R-T
HCJ8	2b	R-T
CH610	2c	R-T
NE92	2d	R-T
BNL3	2e	R-T
FR4	2f	P-
HCVTR	3b	LRQTLNVV-
DK13	4d	R-TM
CAM600	4e	R-TM
GB809	4e	L-R-TM
BNL7	4 k	R-TM
BE95	5a	R-TM
HK2	6a	LR-TT
FR1	7a	LR-TM
VN4 VN13	8a 8b	LR-TI
VN12	9a	LR-TM
NE98	10a	LR-TXVV-

Figure 2 - continued

HCV1	la	51 100 KTSERSQPRGRRQPIPKARRPEGRTWAQPGYPWPLYGNEGCGWAGWLLSP
HCV-J	1b	W. V. C.
BNL1	1d	X-XSX-XXXXXX
BNL2	1d	
CAM1078	le 1f	AA
FR2	11	•
нсј6	2a	LLL
HCJ8	2b	DST-KS-GK
CH610	2c	LL
NE92	2d	LL
BNL3	2e	
FR4	2f	LL
HCVTR	3b	KQ-HLSRSKKL
DK13	4d	QLS
CAM600	4e	
GB809	4e	
BNL7	4 k	XX
BE95	5a	AL
HK2	6a	Q-QH
FR1	7a	V-Q-TS-G
VN4	8a	V-HQT
VN13	8b	V-HQT
41170	02	
VN12	9a	AV-QNQ
NE98	10a	SRTS

Figure 2 - continued

9420 2	00	
HCV1 HCV-J BNL1 BNL2 FR2	la 1b 1d 1d 1f	101 150 RGSRPSWGPTDPRRRSRNLGKVIDTLTCGFADLMGYIPLVGAPLGGAARANNNS-T
HC-J6 HC-J8 CH610 NE92 BNL3 FR4	2a 2b 2c 2d 2e 2f	VV
HCV-TR	3b	NF
GB116 DK13 CAM600 GB809 G22 GB549 GB438 BNL7	4c 4d 4e 4f 4g 4h 4k	VV -X-XNXVV NVV VV VV
BE95	5a	NNKK
нк2	6a	HNVVVV
FR1	7a	NNXXLVL-GV-A-
VN4 VN13	8a 8b	NNXXXIE
VN12	9a	D-X-NXV-AE
NE98	10a	N

Figure :	2	-	continued
----------	---	---	-----------

HCV1 HCV-J BNL1	1a 1b 1d	THIS WAS A VOICE OF THE TAIL O
BNL2	1d	
FR2	1f	-XXGXXXXXYXZ
	11	-XXGXXXXXXXXXTT-HEAS-V
HC-J6	2-	" THE HST-DG
HC-J8	2a	I-T-VAE-K-ISTG
CH610	2b	I-T-VAE-K-ISTG
	2c	IISSS
NE92	2d	I
BNL3	2e	XIV-GLK-TSSS
FR4	2f	XIXXIV-GLK-TSSS IXXVV-XVE-K-TSQA IIV-IK-NSHF
BNL4	2q	
BNL5	2h	
BNL6	2i	IVVK-TSTM
0	21	IVK-TSHS
HCV-TR	3b	1 VVA-RS-S
ncv-1K	3D	A-GFFCGLEYT-TS
GB116	4 -	CGLEYT-TS
DK13	4c	-EAVISTVNYAS-V
	4d	TVNYAS-V
CAM600	4e	AVITVNYAS-I
GB809	4e	AVITVNYAS-I
G22	4 f	GVNYAS-V
GB549	4 q	AVI
GB438	4h	AV T
BNL7	4 k	
BNL8	4 k	
BNL9	4 k	
BNL9	4 k	
BNL10		
BNL11	4 k	
DIATIT	41	IITNYVS-I
חחסכ	_	
BE95	5a	VPYAS-I

HK2	6a	AII
		AIITTTYGS
FR1	7a	AIIK-AS-I
		AII-K-AS-I
VN4	8a	VVT V
	Ju	XXIXXXXXXXTAHYT-KS
VN12	9a	· • • • • • • • • • • • • • • • • • • •
	Ja	-XAIIXTLNYA-KS
NE98	3.0	I IIIIA-KS
115 30	10a	I-FFFLT-TAGLEYAS

Figure 2 - continued

HCV-1 HCV-J BNL1 BNL2 FR2	la lb ld ld lf	250 YHVTNDCPNSSIVYEAADAILHTPGCVPCVREGNASRCWVAMTPTVATRDSM-MS-FLL-A-NSIMDGM-M-YD-HLM-LL-VKXLSIMSGMAN-SMXLL-VK-
HC-J6 HC-J8 CH610 NE92 BNL3 FR4 BNL4 BNL5 BNL6	2a 2b 2c 2d 2e 2f 2g 2h 2i	-MT-DTWQLQA-VVEKVTIPVS-NVQQ -YAS-NTWQLTVLENDNGTLHIQVNVKH -MSWQLEG-VEQIPVS-NI-Q -MQWQLRVVEEKIIPVS-NI-VSQ -MAS-NWQLXVVENSSGRFHIPIS-NI-VSK -MAA-DWQLRVVE-SRTFT-VS-NVSR -MAS-NIWQMQG-VVELQKIPVNVNQ -MS
HCVTR	3b	-VLS-GE-VLTTQ-STTVSTV-T
GB116 DK13 CAM600 GB809 G22 GB549 GB438 BNL7 BNL8 BNL9 BNL9 BNL10 BNL11 GB724	4c 4d 4e 4f 4f 4h 4k 4k 4k 4k	I
BE95	5 a	DNLAMTVQILSAPS
HK2	6a	LLDAMLLVDDR-TH-VL-IPN
FR1	7a	LS-NFETMLIKAELPVSL-VPN
VN4	8a	LQASL-VPN
VN12	9a	LNGMLKTLTKLSASL-VQN
NE98	10a	-MS-GG-ILSTIPVSXVKS

Figure 2 - continued

		251
HCV-1	la	
HCV-J	lb	GKLPATOLRRHIDLLVGSATLCSALYVGDLCGSVFLVGQLFTFSPRRHWT
BNL1	1b	SSI-T-TIVA-AMYE-
BNL2		ASV-TXAIVXX-FM-XAM-H-
	1d	ANV-TAAIVT-AFRMLYH-
FR2	1f	ANA-IDEVVA-VFM-IGTS
HC-J6	2a	PGALTQGTMV-MG-M-AA-M-TVOHF
HC-J8	2b	RGALTRST-V-MI-MAAVA-MTIC-A-MVO-NE
CH610	2c	PGTLTKGA-V-VI-MVDIMIDD-DVID-O
NE92	2d	PGALTKG-TTIIAFIA-M-AS-V-IIOH-KF
BNL3	2e	PGALTKGARAV-MVA-MIAA-A-IVA-KYF
FR4	2f	PGALTRGATI-MIA-MIAA-VAVVQY-TF
BNL4	2g	PGALTRG-TTI-MVIV-A-MIAA-VVIVOH-NF
BNL5	2h	PGALTRG-TTI-AVF-A-M-S-F-MIOH-IF
BNL6	2i	PGAXTKG-TII-AF
DIADO	21	EGAVIVGIII-YK
HCVTR	3b	I CIMMINOT M II W DDA
HCVIK	30	LGVTTASI-T-V-MARQAF-AART-
CD116		••••
GB116	4c	VGA-LESS-VMAVIGM-S-Q
DK13	4 d	LNA-LESVMGIVG
CAM600	4e	AGA-LEPVMAMIGIMO
GB809	4e	VGA-LEPVMAVGIMO
G22	4f	LGA-LESMVMT
GB549	4 q	VGA-LESMVMAVIGMRE
GB438	4ñ	LGA-L-SV-Q-V-M-AI-H-GA-MVS-Q
BNL7	4 k	IGA-LESS-VMAVIX-XGLM-S-R
BNL8	4 k	IGA-LESS-VM-AVIGLM-S-R
BNL9	4 k	IGA-LESS-VMAVIGAM-S-R
BNL9	4 k	TAA-LESS-VMAVI-XGLM-SXQ
BNL10	4 k	TCA-IFCC-V-TM A TVT-XGLM-SXQ
BNL11	41	IGA-LESS-V-VMAVIGLM-S-R
GB724		LSA-LMSVV-MASGAMQ
GD / 24	4 x	VDA-LESFVMAVGAMQ
DD05	-	
BE95	5a	LGAVTAPAV-Y-A-G-AAALMYRO-A-
HK2	6a	ASTGFVA-A-VVSILAQ
		· · · · · ·
FR1	7a	SSV-IHGFVA-AFM-IIIR-KY-OV
		11 11 11 11 11 1 1 1 1 1 1 1 1 1 1 1 1
VN4	8a	AST-V-GF-K-V-IMA-AFMGLLRM-OV
		191 1 01 1 1 11 11 11 M
VN12	9a	ASVSIRGV-E-VA-AFMGLRMYEI
	24	TO A DITTO A D
NE98	10a	DCNAMAC M. 17 AM VA
-14-70	IVa	PCAATAST-V-MM-XAALXG-SWRH-Q

Figure 2 - continue	ri mire	2	_	continue	d
---------------------	---------	---	---	----------	---

HCV-1 HCV-J BNL1 BNL2 FR2	la 1b 1d 1d	301 319 TQGCNCSIYPGHITGHRMA V-DVSE V-DSXXX
HC-J6 HC-J8 CH610 NE92 BNL3 FR4 BNL4 BNL5	2a 2b 2c 2d 2e 2f 2g 2h	V-DX V-EX V-DX V-E
HCVTR	3b	V-TVS
GB116 DK13 CAM600 GB809 G22 GB549 GB438 BNL7 BNL8 BNL9 BNL9 BNL9 BNL10 BNL11 GB724	4c 4d 4e 4f 4g 4h 4k 4k 4k 4k	DAVDTDTD
BE95	5a	V-NSV
HK2	6a	V-DTV
FR1	7a	DXNXV
VN4	8a	V-ET
VN12	9a	A-DA
NE98	10a	V-D

Figure 3

SEQ ID NO. 1 (BNL1, 1d)
ATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGTAACACCAACCGCCGCCCTCAKGGSGTN
NNNNNNCCGGGTGGCGGTCAGATCGTTGGTGGAGTTTACCTGTTGCCGCGCAGGGGCCCCCAGGNNG
GGTGTGCGCGCGACTAGGAAGACTTCCGAGCGGTCACAACCTCGTGGCAGGCGACAGCCTATCCCC
AAGGCTCGYCGGYCCGAGGGCAGGTCCTGGGCTCAGCCCGGGTATCCTTGGCCCCTCTATGGCAAT
GAGGGCTGCGGGTGGGCGGGTTGGCTCCTGTCCCCCCCGCGGCTCTCGGCCCAATTGGGGCCCC

SEQ ID NO. 3 (BNL1, 1d)
GACGGCGTGAACTATGCAACAGGGAACTTGCCCGGTTGCTCTTTCTCTATCTTCCTCTTTGGCTTTG
CTGTCCTGCTTGACGGTTCCAACKACCGCTCACGAGGTGCGCAACGCATCCGGGGTGTATCATGTC
ACCAACGACTGTTCCAACTCGAGCATCATCTATGAGATGGACGGTATGATCATGCACTACCCAGGG
TGCGTGCCCTGCGTTCGGGAGGATAACCATCTCCGCTGCTGGATGGCGCTCACCCCCACGCTTGCG
GTCAAAAAYGCTAGTGTCCCCACTRCGGCAATCCGACGTCACGTCGACTTGCTTGTTGGGGGNNCC
ACGTTCTGTTCCGCTATGTACGTGGGRGACCTTTGCGGGGTCTCTCTCCTCGCTGGCCAGCTATTC
ACCTTTTCACCCCGCATGCACCATACAACGCAGGAGTGCAACTGCTCAATC

SEQ ID NO. 5 (BNL2, 1d)
ATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGTAACACCAACCGCCGCCCACAGGACGTC
AAGNTCCCGGGTGGTCGTCGTTGGTGGAGTTTACCTGTTGCCGCGCAGGGGCCCCAGGTTG
GGTGTGCGCGCGACCAGGAAGACTTCCGAGCGGTCGCAGCCTCGTGACAGGCGACAGCCTATTCCT
AAGGCTCGCCAGTCCGATGGCAGNNCCTGGGCTCAGCCAGGCCATCCCTGGCCCCTCTATGGCAAT
GAGGGCTGCGGATGGGCGGGTCCCTGTCCCCCCGCGGCTCTCGGCCCAGTTGGGGCCCC

SEQ ID NO. 7 (BNL2, 1d)
GACGGCGTGAACTATGCAACAGGGAATTTGCCTGGTTGCTCTTTCTCTATCTTCCTCTTAGCTTTT
CTGTCCTGCTTGACGGTTCCAACTACCGCTCATGAGGTGCGCAACGCATCCGGGGTATATCATCTC
ACCAATGACTGTTCCAACTCGAGCATCATCTATGAGATGAGTGGTATGATCTTGCACGCCCCAGGG
TGTGTGCCCTGCGTTCGGGAGAACAACTCTTCTCGTTGCTGGATGCCRCTCACCCCCACGCTTGCG
GTCAAAGACGCTAATGTCCCTACTGCGGCAATCCGACGCCATGTCGACTTGCTGGTTGGGACAGCC
GCGTTTCGTTCCGCTATGTACGTGGGGGACCTCTGCGGATCCGTCTTCCTTGTCGGCCAGCTATTC
ACCTTTTCACCCCGCTTGTACCATACAACACAGGAGTGCAACTGCTCAATC

Figure 3 - continued

SEQ ID NO. 11 (FR2, 1f) ATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGCAACACCAACCGCCGCCCACAGGACGTT AAATTCCCGGGTGGGGGCAGATCGTGGGTGGAGTTTACTTGTTGCCGCGCAGGGGCCCCCAGGTTG GGTGTGCGCGCGACGAGGAAGACTTCCGAGCGGTCGCAACCTCGCGGAAGGC GACAGCCTATCCCCAAGGCTCGCCGACCCGAGGGCAGGTCCTGGGCTCAGCCTGGGTACC CATGGCCCCTCTATGCTAACGAGGGCTGCGGATGGGCGGGATGGCTCCTGTCCCCTCGCG GCTCCCGTCCTAGCTGGGGCCCCAATGACCCCCGACGTAGATCACGCAATTTGGGTAAGG TCATCGATACCCTAACGTGTGGCTTCGCCGATCTCATGGGGTACATTCCGCTCGTCGGCGC CCCCCTAGGGGGCGCTTCCAGAACCCTGNCACATGGTGTCCGGGTCCTGGNAGGCGGCGTGATNNN NNNNNNNNAACCTTCCNGGTTGCTCTTTNNCTATCTTCCTCTTGGCNTTACTCTCTTGCCTCAC AGTCCCCACCTCTGCCTATGAGGTGCACAGCACAACCGATGGCTACCATGTCACTAATGACTGTTC CAACGCCAGCATCGTATATGAGGCAAAGGACATCATCCTTCACACGCCTGGGTGNGTGCCCTGCAT ACGGGAAGGCAATATCTCCCGTTGCTGGGTACCGCTCACCCCCACGCTCGCAGCGCGGATCGCGAA CGCTCCCATCGATGAGGTGCGGCGTCACGTCGACCTCCTCGTGGGGGCAGCCGTGTTCTGCTCAGC CATGTACATTGGGGACCTTTGTGGGGGGCGTCTTCCTCGTTGGGCAATTGTTCACCTTCACGTCCCG GCGGCATTGGACGGTGCAGGACTGTAATTGTTCCATTTACTCTGGCCACATAACGGGCCACCGNNN NNNN

SEQ ID NO. 13 (BNL3, 2e)
ATGAGCACAAATCCTAAACCTCAAAGAAAAACCAAAAGAAATACCAACCGCCGCCCACAGGACGTC
AAGTTCCCGGGCGGCGGCCAGATCGTTGGCGGAGTTTACTTGTTGCCGCGCAGGGGCCCCAGATTG
GGTGTGCGCGCGACGAGAAAGACTTCTGAACGGTCCCAGCCACGTGGAAGGCGCCCAGCCCATCCCT
AAAGATCGGNGNGCCACTGGCAGGTCCTGGGGACGTCCAGGATATCCCTGGCCCCTGTATGGGAAC
GAGGGGCTCGGCTGGGCAGGATGGCTCCTGTCCCCCCGAGGCTCTC

SEO ID NO. 17 (FR4, 2f) ATGAGCACAAATCCTAAACCTCAAAGAAAAACTAAAAGAAACACTAACCGTCGCCCACAGGAC GTTAAGTTCCCGGGCGGCGGCCAGATCGTTGGCGGAGTTTACTTGTTGCCGCGCAGGGGCCCCAG GTTGGGTGTGCGCGCCAAGGAAGACTTCTGAACGGTCCCAGCCACGTGGAAGGCGCCAGCCC ATCCCAAAAGATCGGCGCCCCCTGGCAAGTCCTGGGGACGTCCAGGATACCCTTGGCCCCTGT ACGGGAACGAGGGCCTCGGCTGGGCAGGGTGGCTCCTGTCCCCCCGGGGGCTCTCGCCCCTCGTG GGGCCCAAACGACCCCCGGCACAGGTCACGCAACTTGGGTAAGGTCATCGATACCCTCACGTG TGGCTTTGSCGACCTCATGGGGTACATACCTGTCGTCGGCGCCCCTGTGGGCGGCGTTGCCAGA GCCCTCGCGCATGGCGTGCGGGTCCTGGAGGACGGGATAAATTATGCAACAGGGAACTTGCCCGGT GTTAAGAACAACAGCCACTTCTACATGGCGACTAATGACTGTGCCAATGACAGCATCGTCTGGCAG CTCAGGGACGCGGTGCTCCATGTTCCTGGATGTGTCCCCTGTGAGAGGTCAGGTAATAGGACCTTC TGTTGGACAGCGGTCTCGCCCAACGTGGCTGTGAGCCGACCTGGTGCTCTCACTAGAGGTCTGCGG GCTCACATTGATACCATCGTGATGTCCGCCACCCTCTGCTCTGCCCTATACATAGGGGACCTATGC GGCGCTGTGATGATAGCAGCGCAAGTTGCCGTCGTCTCACCGCAATACCATACTTTTGTCCAGGAA TGCAACTGCTCCATATACCCAGGCCATATCACAGGACATCGAATGGNN

Figure 3 - continued

SEQ ID NO. 19 (BNL4, 2g)
GACGGGGTAAATTATGCAACAGGGAATCTGCCTGGTTGCTCTTTCTCTATCTTCTTGTTGGCTCTT
CTGTCTTGTGTCACCGTGCCTGTCTCTGCCGTGCAGGTTAAGAACACCAGTACCATGTACATGGCA
ACCAATGACTGTTCCAACAACAGCATCATCTGGCAAATGCAGGGCGCGGTGCTTCATGTTCCTGGA
TGTGTCCCGTGTGAGTTGCAGGGCAATAAGTCCCGGTGCTGGATACCGGTCACTCCCAACGTGGCT
GTGAACCAGCCCGGCGCCCTCACTAGGGGCTTGCGGACGCACATTGACACCATCGTGATGGTCGCT
ACGCTCTGTTCTGCACTCTACATCGGGGACGTGTGTGGCGGGTGATGATAGCTGCTCAGGTTGTC
ATTGTCTCGCCGCAACATCACAACTTTTCCCAGGATTGCAATTGTTCCATC

SEQ ID NO. 21 (BNL5, 2h)
ATGAGCACAAATCCTAAACCTCAAAGAAAAACCAAAAGAAACACTAACCGCCGCCCACAGGACGTT
AAGTTCCCGGGCGGTGGCCAGATCGTTGGCGGAGTATACTTGTTGCCGCGCAGGGGCCCCCGGTTG
GGTGTGCGCGCGACGAGAAAACTTCCGAACGGTCCCAGCCACGTGGGAGGCCCCAGCCCATCCCT
AAAGATCGGCGCTCCACTGGCAAATCCTGGGGACGTCCAGGATACCCTTGGCCCCTGTATGGGAAC
GAGGGCCTTGGTTGGGCAGGATGGCTCTTGTCCCCTCGAGGCTCTC

SEQ ID NO. 23 (BNL5, 2h)
GACGGGATAAACTACGCAACAGGGAATCTGCCCGGTTGCTCCTTTTCTATCTTCTTGCTGGCCTTG
CTATCCTGTCTCACTGTGCCGGCGTCCGCTGTGCAGGTCAAGAACACCAGCCACTCTTATATGGTG
ACCAATGATTGCTCAAACAGCAGCATTGTCTGGCAGCTTAAAGGATGCTGTGCTTCACGTCCCTGGA
TGTGTTCCATGTGAGAGGCACCAAAATCAGTCTCGCTGCTGGATACCTGTGACACCCAATGTGGCC
GTGAGCCAACCTGGCGCGCTCACCAGGGGTTTGCGGACGCACATTGACACCATCGTTGCGTCTGCT
ACCGTCTGCTCAGCTTTGTATGTGGGCGACTTCTGCGGCGCGCAGTGATGTTGGTCTCTCAATTTTTC
ATGATCTCCCCTCAGCACCACATCTTCGTCCAGGATTGCAACTGCTCGATA

SEQ ID NO. 27 (BNL7, 4k)
ATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGTAACACCAACCGCCGCCCCCATGGACGTT
AAGTTCCCGGGTGGTGGCCAGATCGTTGGCGGAGTTTACTTGTTGCCGCGCAGGGGCCCCAGGTTG
GGTGTGCGCGCGACTCGGAAGACTTCGGAGCGGTCGCAACCTCGTGGGAGACGCCAACCTATCCCC
AAGGCGCGTCGATCCGAGGGAAGGTCCTGGGCACAGCCAGGATATCCATGGCCTCTTTACGGTAAT
GAGGGTTGCGGGTGGGCANNATGGCTCTTGTCCCCCCGCGGTTCTC

SEQ ID NO. 29 (BNL7, 4k)
GACGGGATCAATTTTGCAACAGGGAACCTCCCCGGTTGCTCCTTTTCTATCTTCTCTTTGGCACTC
CTCTCGTGCCTGACTGTCCCCGCTTCGGCCATCAACTATCGCAATGTCTCGGGCATTTACTATGTC
ACCAATGATTGCCCGAATTCAAGCATAGTGTATGAGGCCGACCATCACATCTTGCACCTCCCAGGT
TGCGTGCCCTGCGTGAGAGAGGGGGAATCAGTCACGTTGCTGGGTAGCCCTTACCCCTACCGTCGCA
GCGCCATACATCGGCGCCCACTTGAGTCTCTACGGAGTCATGTGGACTTGATGGTGGGGGCCGCC
ACTGTTTGTTCAGCCCTTTACATCGGGGATTTRTGTGGYGGCTTGTTCCTAGTCGGTCAGATGTTC
TCTTTCCGACCAAGGCGCCACTGGACTACTCAAGATTGCAATTGTTCCATC

Figure 3 - continued

SEQ ID NO 31 (BNL8, 4k)

SEQ ID NO. 33 (BNL9, 4k)

SEQ ID NO. 35 (BNL10, 4k)

GACGGGATCAATTATGCAACAGGGAATATTCCCGGTTGCTCYTTTTCTATCTTCCTTYTGGCACTT
CTCTCGTGTCTGACTGTCCCCGCTTCGGCCACTAACTATCGCAACGTCTCGGGCATCTACCATGTC
ACCAATGACTGCCCGAATTCAAGCATAGTGTATGAGGCCGACCATCACATCTTAGCACTTCCAGGT
TGCGTGCCCTGCGTGAGAGTGGGGAACCAGTCACGCTGCTGGGTGGCCCTTACCCCTACCGTCGCA
GCGCCATACACCGCGGCGCCGCTTGAGTCCCTGCGGAGTCATGTGGATCTTGATGGTGGGAGCTGCC
ACTGTTTGTTCAGCCCTTTACATCGGGGAYTTGTGTGGCGGCTTGTTCTTGGTTGGTCAGATGTTC
TCTTTYCAGCCTCGGCGCCACTGGACTACCCCAGGATTGCAATTGTTCCATC

SEQ ID NO. 37 (BNL11, 4k)

SEO ID NO. 39 (BNL12, 41)

SEQ ID NO. 45 (VN13, 7a)

Figure 3 - continued

SEQ ID NO. 43 (VN4, 7c) ATGAGCACACTTCCAAAACCCCAAAGAAAAACCAAAAGAAACACCATCCGCCGCCCACA GGACGTCAAGTTCCCGGGTGGCGGCCAGATCGTTGGTGGAGTCTACTTGCTGCCGCGCAG GGGCCCGCGCTTGGGTGTGCGCGCGACGAGAAAGACTTCTGAACGGTCCCAGCCCAGAGG TAGGCGCCAACCAATACCCAAAGTGCGCCACCAAACGGGCCGTACCTGGGCCCAGCCCGG CCGCGGCTCTCGCCCAAATTGGGGCCCAAACGACCCCCGGCGGAGGTCCCGCAACTTGGG TAAAGTCATCGACACCCTTACTTGCGGCTTCGCCGACCTCATGGGGTATATCCCTGTCGTAG GCGCTCCGWTGGGAGGCGTCGCGGNGGCCTTGGCGCATGGGGTCANGGNCATCGAGGACGGNGTAA ATTACGCAACAGNGAATCTTCCCGGNNGCTCTNTCTCTATCTTNCTCTTGGCACTTCTCTCGTGCC TTACAACACCAGCCTCCGCGGCGCATTATACCAACAAGTCTGGCCTGTACCATCTCACCAACGACT GCCCCAACAGCAGCATCGTTTATGAGGCGGAGACACTGATTTTGCACTTGCCTGGGTGTGTACCTT GTGTGAAGRTGRACAATCAATCCCGGTGCTGGGTGCAGGCCTCCCCGACCCTGGCAGTGCCGAACG CGTCTACGCCAGTCACCGGGTTCCGCAAACATGTGGACATCATGGTGGGCGCTGCCGCGTTCTGTT CAGCTATGTATGTGGGGGACCTGTGCGGGGGCCTTTTCCTCGTTGGACAGCTCTTCACGCTCAGGC CTCGGATGCATCAGGTTGTCCAGGAGTGTAACTGTTCCATCTACACAGGGCATATCACTGGACACC GAATGGCA

SEQ ID NO. 47 (VN12, 7d)

SEQ ID NO. 41 (FR1, 9a) ATGAGCACACTTCCAAAACCCCAAAGAAAAACCAAAAGAAATACTAACCGTCGCCCTATGGAC GTCAAGTTCCCGGGCGGCGGCCAGATCGTTGGTGGAGTTTACTTGTTGCCGCGCAGGGGC CCTCGTTTGGGTGTGCGCGCGACGAGAAAGACCTCCGAACGGTCCCAGCCTAGAGGCAGG CGCCAGCCCATACCAAAGGTACGCCAGCCGACAGGCCGTAGCTGGGGTCAACCCGGCTAC CCTTGGCCCCTTTATGGCAACGAGGGCTGCGGATGGCGGGATGGCTCCTGTCCCCCGC GGGTCTCGTCCTAATTGGGGCCCCAACGACCCCCGGCGAAGGTCCCGCAACTTGGGTAAG GTCATCGATACCCTTACATNCGGNCTAGCCGACCTCATGGGGTACATCCCTGTCCTAGGAGG GCCGCTTGGCGCGTTGCGGCTGCCCTGGCGCATGGCGTTAGGGCAATCGAGGACGGGGTCAATTA CGCAACAGGGAATCTTCCTGGTTGCTCCTTTTCTATCTTCCTCTTAGCACTGTTATCGTGCCTCAC TACACCAGCCTCAGCAATTCAAGTCAAGAACGCCTCTGGGATCTACCATCTTACCAATGACTGCTC GAACAACAGCATCGTTTTTGAGGCGGAGACCATGATACTGCATCTTCCAGGTTGTGTCCCATGTAT CAAGGCGGGGAATGAGTCACGATGTTGGCTCCCTGTCTCCCCCACCTTAGCCGTCCCCAACTCATC AGTGCCAATCCACGGGTTTCGCCGACACGTAGACCTCCTCGTTGGGGCAGCGGCATTTTGTTCGGC CATGTACATCGGAGACCTCTGTGGTAGCATAATCTTGGTAGGGCAGCTTTTTACTTTCAGGCCTAA GTACCATCAGGTTACCCAGGATTGTAACTGCTCTATNAACNCTGGCCACGTCACGGGACACAGGAT GGCA

Figure 3 - continued

SEQ ID NO. 49 (NE98, 10a)

SEQ ID NO. 51 (NE98, 10a)

SEQ ID NO. 53 (BNL1,1d)

SEQ ID NO. 55 (BNL2,1d)

CTCGACAGTTACTGAGAACGACATCCGTACCGAGGRATCAATCTATCAATGTTGTGACTTGGCCCC
YGAGGCCCGCAAGGCCATAAAGTCGCTCACCGAGCGGCTGTACGTCGGGGGCCCCCTAACCAATTC
AAAGGGGCAGAACTGCGGCTATCGTCGGTGTCGCGCTAGCGGCTGCTGACCACCAGCTGCGGCAA
CACCCTCACATGCTACTTGAAAGCCAGGGCGGCCTGTCGAGCTGCAAAGCTCCAGGACTGCACGAT
GCTCGTGTGCGGAGACCACCTTGTCGTTATCTGTGAGAGCGCGGGAGTCGAGGAGGACGCGGCGAA
CCTACGAGTC

SEQ ID NO. 57 (FR17,1d)

SEQ ID NO. 59 (CAM1078,1e)

CGTACAGCCTCCAGGACCCCCCTCCCGGGAGAGCCATAGTGGTCTGCGGAACCGGTGAG
TACACCGGAATTGCCAGGACGACCGGGTCCTTTCTTGGATCAACCCGCTCAATGCCTGGA
GATTTGGGCGTGCCCCCGCAAGACTGCTAGCCGAGTAGTGTTGGGTCGCGAAAGGCCTTG
TGGTACTGCCTGATAGGGTGCTTGCGAGTGCCCCGGGAGGTCTCGTAGACCGTGCACCAT
GAGCACGAATCCTAAACCTCAAAGAAAAACCAAAAGAAACACCAACCGCCGCCCACAGGA
CGTCAAGTTCCCGGGCGGTGGCCAGATCGTTGGTGGAGTCTACCGCGCAGCGG
CCCTAGATTGGGTGCGCGCGCGGAAGACTTCGGAGCGGTCGCAACCTCGTGGGAG
GCGCCAACCTATTCCCAAGGAGCGCCGACCCGAGGGCAGGTCCTGGGCCAGCCCGGGTA
CCCCTGGCCCCTCTATGGTAACGAGGGCTGCGGGTGGGCAGGTCCTGTCCCCTCG
CGGCTCCCGTCCTAGTTGGGGTCCTACTGACCCCCGGCGTAGGTCACCGCAATTTGGGTAA
GGTCATCGATACCCTCACGTGTTGNTTCGCCGACCTCATGGGGTACATACCG

Figure 3 - continued

SEO ID NO. 61 (CAM1078, 1e)

CTCAACGGTCACTGAAGCTGATATCCGAACAGAGGAGTCCATATACCAATGCTGTGACCTGCACCC CGAAGCACGTGTAGCCATCAAGTCTTTGACTGAAAGGCTGTACGTCGGGGGGCCCTTGACCAATTC AAAAGGGGAGAACTGCGGCTATCGCAGATGCCGTGCCAGCGGCGTCTTGACAACCAGCTGCGGCAA CACCCTCACCTGCTATATCAAGGCCCTAGCAGCCTGTAGAGCTGCCAAGCTCCAGGACTGCACCAT GCTCGTCTGTGGCGACGACCTGGTCGTGATCTGCGAGAGTGTAGGGACCCAGGAGGATGCGGCGAG CCTGCGAGCC

SEO ID NO. 63 (FR2, 1f)

NTCAACAGTCACTGAGAGTGATATCCGTACAGAGGAGTCCATCTACCAATGCTGTGATCTAGACCC CGAGGCTCGCAAGGCCATAAGGTCCCTCACAGAGGGCTTTATATCGGGGGTCCCCTGACAAACTC AAAAGGGCAGAACTGCGGCTACCGCCGATGCCGTGCAAGCGGCGTCCTGACGACTAGCTGCGGCAA CACCCTCACCTGTTACATAAAGGCCAGGGCAGCCTGTCGAGCTGCGAAGCTCCAGGATTGCTCAAT GCTCGTCTGTGGCGACCCTTGTCGTTATCTGCGAGATCGAGGGGTCCANGAGGATCCGTCGAN NNNNNNNNNN

SEO ID NO. 65 (FR16,19)

SEO ID NO. 67 (FR16,1g)

SEO ID NO. 69 (BNL3, 2e)

CTCGACAGTCACAGAGAGAGATATAAGNACTGAGGAGTCCATATACCAGGCTTGTTCCTTACCCGA GCAGGCCAGAACTGCCATACACTCATTGACTGAGAGACTCTACGTAGGAGGGCCCATGATGAACAG CAAAGGGCAATCCTGCGGATACAGGCATTGCCGCGCCAGCGGAGTGCTCACCACCAGTATGGGGAA TACCATCACGTGCTACATCAAGGCCCTAGCGGCTTGTAAAGCAGCAGGAATAGTGGCCCCCACCAT GCTGGTGTGCGGCGATGACCTAGTTGTCATCTCAGAGAGTCAGGGAGTCGAGGAGGACGACCGGAA CCTGANNNNN

SEQ ID NO. 71 (FR4, 2f)

CTCAACCGTCACAGAGAGGGATATAAGAACTGAGGAGTCCATATACCTGGCCTGCTCCTTACCCGA GCAGGCCCGGACTGCCATACATTCATTAACTGAGAGACTTTACGTGGGAGGGCCCATGATGAACAG CAAAGGGCAGTCCTGCGGATACAGGCGTTGCCGCGCTAGCGGAGTGCTCACCACCAGTATGGGGAA CACCATCACGTGTTATGTGAAAGCCCTCGCAGCTTGTAAAGCTGCGGGCATTGTTGCCCCCACGAT GCTGGTGTGCGGCGATGACCTGGTTGTCATCTCAGAGAGTCAGGGGGCTGAGGAGGACGAGCGAAA CCTGAGAGTC

SEQ ID NO. 73 (BNL5,2h)

CTCAACAGTCGCGGAGAGACATCAGGACCGAGGAGTCCATTTACCTTGCCTGCTCCTTACCCGA GCAAGCCCGAACTGCCATACATTCATTGACTGAGAGACTTTACGTAGGAGGGCCCATGATGAACAG CAAGGGACAGTCCTGCGGTTACAGACGTTGCCGCGCCAGCGGAGTGCTCACCACCAGCATGGGGAA TACCATCACATGCTATGTGAAGGCATTAGCTGCCTGCAAAGCTGCAGGCATCGTTGCTCCCACGAT GCTGGTTTGTGGCGACGATCTGGTCATCATCTCAGAGAGTCAGGGAACCGAGGAGGATGAGCGGAA CCTGAGAGTC

SEQ ID NO. 75 (FR13, 2k)

CGNACANCCTCCAGGCCCCCCCCCCCCGGGAGAGCCATAGTGGTCTGCGGAACCGGTGAG TACACCGGAATTGCCGGGAAGACTGGGTCCTTTCTTGGATAAACCCACTCTATGCCCGGC CATTTGGGCGTGCCCCCGCAAGACTGCTARCCGAGTAGCGTTGGGTTGCGAAAGGCCTTG TGGTACTGCCTGATAGGGTGCTTGCGAGTGCCCCGGGAGGTCTCGTAGACCGTGCATCAT GAGCACAAATCCTAAACCTCAAAGAAAAACCAAAAGAAACACTAACCGCCGCCCACAGGA CGTTAAGTTCCCGGGCGGTGGCCAGATCGTTGGCGGAGTATACTTGTTGCCNTGCAGGGG NCCCAGGTNGNGTNTATGCGCAACGANGAAGACTNCCGAACAGTCCCAGCCACGTGGGAG GCGCCAGCCCATCCCGAAAGATCGGNGCACCACTGGCAAGTCCTGGGGACGTCCAGGATA TCCCTGGCCCCTGTATGGGAACGAGGGCCTCGGGTGGGCAGGGTGGCTCCTGTCCCCCG GGGCTCCCGCCCGTCATGGGGCCCCACGGACCCCCGGCATAGGTCGCGCAACTTGGGTAA GGTCATCGATACCCTCACGTNCGGCTTTNCCGACCTCATGGGGTACATTCCCGTCGTTGG CGCCCCAGTAGGNGGCGTCGCCAGAGCTCTCGCGCATGGCGTGAGAGTCCTGGAGGACGG TCTGTCCTGAATTACCGNGCCAGTTTCTGCTGTGGAAATCAAAAACACCAGMAACACATA CATGGTGACTAACGACTGTTCAAACAGYAGCATCACCTGGCAGCTTNNGNNCGCGGTGCT TCACGTTCCTGGATGCGTCCCCTGTGAACGAGGGGCAACAGTTCCCGGTGCTGGATTCC AGTCACGCCCRACGTAKNCGTGAGCCGACCTGGTGCCCTAACCGAGGGTTTGCGATCGCA CATCGACACCATCGTAGCGTCCGCAACATTTTGTTCTGCCCTCTACATAGGGGATGTATG TGGCGCGATAATGATAGCTGCCCAAGTGGTCATCGTCTCGCCGGAGCATCATCACTTTGT CCAGGACTGTAACTGTTCCATCTACCCGGGCCACATAACGGGGCCTCGTATGTNG

SEQ ID NO. 77 (FR13,2k)

ATCCACAGTCACTGAAAGAGACATCAGAGTTGAAGAGTCCGTTTATCTGTCCTGTTCACTTCCCGA GGAGGCCCGAGCTGCCATACACTCACTAACTGAGAGGCTGTACGTGGGAGGTCCCATGCAGAACAG CAAGGGGCAATCCTGCGGATACAGGCGCTGCCGCGCCAGCGGGGTGCTCACCACTAGCATGGGGAA TACTCTCACATGCTACTTGAAGGCCCAGGCGGCCTGCAGGGCCGGGGCATTGTTGCACCCACAAT GCTGGTGTGTGTGGCGACGACCTGGTCGTCATCTCAGAGAGTCAGGGGACTGAGAGGGACGAGAACAA CCTGAGACCT

SEQ ID NO. 79 (FR18,21)

CTCAACAGTCACGGAGAGGGACATCAGGAATGAGGAGTCCATATTCCTGGCCTGCTCGTTGCCCGAGGAGGCCCGGACTGTCATACATTCGCTCACTGAGAGACTCTACATAGGCGGGCCGATGATGAACAGCAAAGGCCAGTCCTGTGGATACAGGCGTTGTCGCGCCAGCGGGGTGTTCACCACTAGCATGGGCAATACCATCACGTGCTATGTGAAAGCCATGGCAGCTTGCAGAGCTGCCGGGATTGACGCCCCCACAATGTTGGTATGTGGCACCTGGTGGTCATCTCAGAGAGTCAGGGGACCGAGGAGGACGAGCGAAATCTGAGAGTC

SEQ ID NO. 81 (PAK64,3g)

CTCTTGACTCTACTGTCACTGAACAGGATATCAGGGTAGAAGAAGAATATACCAATGTTGTGACC
TTGAGCCGGAGGCTAGACGGGCAATCAAATCGCTCACGGAACGGCTTTACGTTGGAGGTCCCATGT
TCAACAGCAAGGGGCTCAAATGCGGATATCGCCGTTGCCGTGCTAGCGGTGTATTGCCCACTAGCT
ACGGTAATACAATCACCTGCTACATCAAGGCCAGAGCGGCTGCTCGAGCTGCGGGCCTTCAAGACC
CATCATTCCTTGTCTGCGGAGATGATTTGGTGGTAGTGGCTGAGAGTTGCGKCGTTGATGAGGAGG
ATAGGGCAGC

SEQ ID NO. 83 (BNL8, 4k)

CTCCACTGTAACCGAAAAGGACATCAGGCCCGAGGAAGAGGTCTATCAGTGTTGTGACCTGGAGCCCGAAGCTCGCAAGGTTATTACCGCCCTCACAGAAAGACTCTACGTGGGCGGCCCCATGCACAACAGCAAGAGAGACCTTTGTGGGTATCGGAGATGCCGCGCAAGCGGCGTCTACACGACCAGCTTCGGAAACACTGACGTGCTACCTCAAAGCCTCAGCTGCTATTAGAGCGGCAGGGCTGAGAGACCCACCATGCTGGTTTGCGGTGACGACTTGGTCGTCATCGCTGAGAGCGGTAGAGAGGAGATAACCGAGCCCCCCCAAGCC

SEQ ID NO. 85 (BNL12,41)

SEQ ID NO. 87 (EG81, 4m)

SEQ ID NO. 89 (VN13,7a)

CTCAACAGTCACAGAGCGCGATGTCCAGACGGAGCATGACATCTACCAGTGCTGTAAGTTGGAGCCCGCAGCACGACCAGCACCATCACATCGCTTACTGACCGATTGTACTNCGGTGGTCCCATGTNTAACTCTAAAGGTCAGGCATGTGGATACCGTAGGTGCAGGGCCAGTGCGTCTTGACCACCATCCTGGCCAATACTCTGACTTGCTACTTGAAAGCTCAGGCGCATGCAGAGCTGCCGGGCTGAAGGACTTTGACATGTTGGTCTTGCTGCGGAGACCTTGTCGTTATTTCGGAGAGTTTGGGGGTCTCGGAGGACACTAGTGCACGAGCT

SEQ ID NO. 91 (VN4,7c)

CTCGACAGTCACCGAGCGCGACATCCRCACCGAGCACGACATCTACCAATGCTGCCAACTTGACCC GGTGGCACGCAAGGCTATTACATCTCTGACTGAGCGGCTGTACTGCGGWGGGCCCATGATGAACTC CCGTGGTCAATCATGTGGATACCGTAGGTGCCGAGCCAGTGGCGTGCTCACCACGAGCTTGGGCAA TACCCTAACATGCTATTTGAAAGCACAAGCAGCGTGTAGGGCAGCAAAGCTCAAAAACTATGACAT GTTAGTCTGCGGAGACGATCTAGTCGTTATCGCGGAGAGTGGAGGAGTCTCTGAGGATGTTGACGC CCTGCGAGCA

SEQ ID NO. 93 (VN12,7d)

CTCCTCCGTCACGGAGCGTGACATCCGCACTGAACACGACATCTATCAGTGCTGCCAATTAGATCC
GGTAGCACGGAAAGCCATTACATCTCTTACTGAGCGGCTGTACTGCGGCGGCCCCATGTACAACTC
TCGAGGTCAGTCATGTGGGTACCGCAGGTGCCGGGCTAGTGGTGTCTTCACCACAAGCTTGGGCAA
CACCATGACATGCTACCTGAAGGCTCAGGCGGCTTGTAGGGCAGCRAAGCTCAAAAACTTTGACAT
GTTGGTCTGCGGAGACGACCTAGTCGTTATTGCTGAGAGCGGAGGAGTCCCTGAGGATGCCGGGGC
CCTGCGAGTC

SEQ ID NO. 95 (FR1, 9a)

ATCCACAGTCACGGGGCGCGACATACGCACAGAACNAGACATTTACCTGTCCTGCCAGCTCGACCC
AGAGGCCCGGAAAGCCATAAAGTCTCTCACTGAGAGGCTCTATGTCGGGGGCCCTATGTACAACTC
AAAAGGGCCAACTCTGTGGTCAACGCCGATGCCGAGCAAGCGGAGTACTCCCCACAAGCATGGGTAA
CACCATCACATGCTTCCTGAAGGCAACCGCCGCTTGCCGAGCAGCCGGCTTTACAGATTATGACAT
GTTGGTCTGCGGAGACGATTTGGTTGTCGTAACTGAGAGTGCTGGAGTCAACGAGGATATCGCTAA
CCTGCGAGCC

SEQ ID NO. 97 (NE98,10a)

SEQ ID NO. 99 (FR14,11a)

SEQ ID NO. 101 (FR15,11a)

38/74

SEQ ID NO. 103 (FR19,11a)

CGTACAGCCTCCAGGACCCCCCTCCCGGGAGAGCCATAGTGGTCTGCGGAACCGGTGAGTACACC
GGAATTGCCGGGAAGACTGGGTCCTTTCTTGGATTAACCCACTCTATGCCCGGAGATTTGGGCGTG
CCCCCGCAAGACTGCTAGCCGAGTAGCGTTGGGTTGCGAAAGGCCTTGTGGTACTGCCTGATAGGG
TGCTTGCGAGTGCCCCGGGAGGTCTCGTAGACCGTGCACCATGAGCACGAATCCTAAAC
ACAAACCAAAAGAAACACCAACCGCCGCCCACAGGACGTTAAGTTCCCGGGCGGTGGCCAGATCGT
TGGCGGGGTGTACTTGTTGCCGCGCAGGGGCCCCAGAGTGGGTGTGCGCGCGACAAAAGACCTC
GGAGCGGTCCCAGCCGCGTGGGAGGCGCCAACCTATCCCCAAGGTTAGGCCGCACCACCGGCCGTT

SEQ ID NO. 105 (FR19,11a)

SEQ ID NO. 2 (BNL1, 1d)

MSTNPKPQRKTKRNTNRRPXXXXXPGGGQIVGGVYLLPRRGPRXGVRATRKTSERSQPRGRRQPIP KAXRXEGRSWAQPGYPWPLYGNEGCGWAXWLLSPRGSRPNWGP

SEQ ID NO. 4 (BNL1, 1d)

DGVNYATGNLPGCSFSIFLLALLSCLTVPXTAHEVRNASGVYHVTNDCSNSSIIYEMDGMIMHYPG CVPCVREDNHLRCWMALTPTLAVKXASVPTXAIRRHVDLLVGXXTFCSAMYVXDLCGSVFLAGQLF TFSPRMHTTQECNCSI

SEQ ID NO. 6 (BNL2, 1d)

MSTNPKPQRKTKRNTNRRPQDVKXPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRDRRQPIP KARQSDGXXWAQPGHPWPLYGNEGCGWAGWLLSPRGSRPSWGP

SEQ ID NO. 8 (BNL2, 1d)

DGVNYATGNLPGCSFSIFLLAFLSCLTVPTTAHEVRNASGVYHLTNDCSNSSIIYEMSGMILHAPG CVPCVRENNSSRCWMXLTPTLAVKDANVPTAAIRRHVDLLVGTAAFRSAMYVGDLCGSVFLVGQLF TFSPRLYHTTQECNCSI

SEQ ID NO. 10 (CAM1078, 1e)

MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYVLPRRGPRLGVRAARKTSERSQPRGRRQPIP KERRPEGR

SEQ ID NO. 12 (FR2, 1f)

MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KARRPEGRSWAQPGYPWPLYANEGCGWAGWLLSPRGSRPSWGPNDPRRRSRNLGKVIDTLTCGFAD LMGYIPLVGAPLGGASRTLXHGVRVLXGGVXXXXXNLXGCSXXIFLLXLLSCLTVPTSAYEVHSTT DGYHVTNDCSNGSIVYEAKDIILHTPGXVPCIREGNISRCWVPLTPTLAARIANAPIDEVRRHVDL LVGAAVFCSAMYIGDLCGGVFLVGQLFTFTSRRHWT VQDCNCSIYSGHITGHXXX

SEQ ID NO. 14 (BNL3, 2e)

MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KDRXATGRSWGRPGYPWPLYGNEGLGWAGWLLSPRGSRPSWG

SEQ ID NO. 16 (BNL3, 2e)

TCXXADLMGYXPVVGAPVGGXARALAXGVRVLEDGINYXTGNLPGCSFSIFXLALLSCVTVPVSXV EVKNTSQAYMATNDCSNNSIVWQLXDAVLHVPGCVPCENSSGRFHCWIPISPNIAVSKPGALTKGL RARIDAVVMSATLCSALYVGDVCGAVMIAAQAFIVAPKRHYFVQECNCSIYPGHITGHRMA

Figure 3 - continued

SEQ ID NO. 18 (FR4, 2f)
MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRAPRKTSERSQPRGRRQPIP
KDRRATGKSWGRPGYPWPLYGNEGLGWAGWLLSPRGSRPSWGPNDPRHRSRNLGKVIDTLTCGFXD
LMGYIPVVGAPVGGVARALAHGVRVLEDGINYATGNLPGCSFSIFLLALLSCITVPVSAIQVKNNS
HFYMATNDCANDSIVWQLRDAVLHVPGCVPCERSGNRTFCWTAVSPNVAVSRPGALTRGLRAHIDT
IVMSATLCSALYIGDLCGAVMIAAQVAVVSPQYHTFVQECNCSIYPGHITGHRMX

SEQ ID NO. 20 (BNL4, 2g)
DGVNYATGNLPGCSFSIFLLALLSCVTVPVSAVQVKNTSTMYMATNDCSNNSIIWQMQGAVLHVPG
CVPCELQGNKSRCWIPVTPNVAVNQPGALTRGLRTHIDTIVMVATLCSALYIGDVCGAVMIAAQVV
IVSPQHHNFSQDCNCSI

SEQ ID NO. 22 (BNL5, 2h)
MSTNPKPQRKTKRNTNRRPQDVKFPGGGRSLAEYTCARRGKLRRSSMG

SEQ ID NO. 24 (BNL5, 2h)
DGINYATGNLPGCSFSIFLLALLSCLTVPASAVQVKNTSHSYMVTNDCSNSSIVWQLKDAVLHVPG
CVPCERHQNQSRCWIPVTPNVAVSQPGALTRGLRTHIDTIVASATVCSALYVGDFCGAVMLVSQFF
MISPQHHIFVQDCNCSI

SEQ ID NO. 26 (BNL6, 2i)
DGINYATGNLPGCSFSIFLLALLSCITVPVSAVQVANRSGSYMVTNDCSNSSIVWQLEEAVLHVPG
CVPCEWKDNTSRCWIPVTPNIAVSQPGAXTKGLRTHIDIIVASATFCSALYV

SEQ ID NO. 28 (BNL7, 4k)
MSTNPKPQRKTKRNTNRRPMDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP
KARRSEGRSWAQPGYPWPLYGNEGCGWAXWLLSPRGSRPSWGPNDPRRRSR

SEQ ID NO. 30 (BNL7, 4k)
DGINFATGNLPGCSFSIFLLALLSCLTVPASAINYRNVSGIYYVTNDCPNSSIVYEADHHILHLPG
CVPCVREGNQSRCWVALTPTVAAPYIGAPLESLRSHVDLMVGAATVCSALYIGDXCXGLFLVGQMF
SFRPRRHWTTQDCNCSI

SEQ ID NO. 32 (BNL8, 4k)
DGINYATGNLPGCSFSIFLLALLSCLTVPASAINYRNTSGIYHVTNDCPNSSIVYEADHHILHLPG
CVPCVRTGNQSRCWVALTPTVAAPYIGAPLESLRSHVDLMVGAATVCSALYIGDLCGGLFLVGQMF
SFRPRRHWTAQDCNCSI

SEQ ID NO. 34 (BNL9, 4k)
DGINYATGNLPGCSFSIFLLALLSCLTVPASAINYHNTSGIYHITNDCPNSSIVYEADHHILHLPG
CVPCVRVGNQSSCWVALTPTIAAPYIGAPLESLRSHVDLMVGAATVCSALYIGDLCGGAFLVGQMF
SFRPRRHWTTQDCNCSI

SEQ ID NO. 36 (BNL10, 4k)
DGINYATGNIPGCXFSIFLXALLSCLTVPASATNYRNVSGIYHVTNDCPNSSIVYEADHHILALPG
CVPCVRVGNQSRCWVALTPTVAAPYTAAPLESLRSHVDLMVGAATVCSALYIGXLCGGLFLVGQMF
SXOPRRHWTTQDCNCSI

SEQ ID NO. 38 (BNL11, 4k)
DGINYATGXLPGCSFSIFLLALLSCLTVPASATNYRNVSGIYHVTNDCPNSSIVFEADHHILHLPG
CVPCVKEGNHSRCWVALTPTVAAPYIGAPLESLRSHVDVMVGAATVCSALYIGDLCGGLFLVGQMF
SFRPRRHWTTQECNCSI

SEQ ID NO. 40 (BNL12, 41)
DGINYATGNLPGCSFSIFILALLSCLTVPASAQHYRNVSGIYHVTNDCPNSSIVYESDHHILHLPG
CVPCVKTGNTSRCWVALTPTVAAPILSAPLMSVRRHVDLMVGAATLSSALYVGDLCGGAFLVGQMF
TFOPRRHWTVQDCNCSI

SUBSTITUTE SHEET (RULE 26)

SEQ ID NO. 46 (VN13, 7a)

MSTLPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KVRHQTGRTWAQPGYPWPLYGNEGCGWAGWLLSPXGSRPNWGPNDPRXRSRNLGKVIDTLTXXFAD LIEYI

SEQ ID NO. 44 (VN4, 7c)

MSTLPKPQRKTKRNTIRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KVRHQTGRTWAQPGYPWPLYGNEGCGWAGWLLSPRGSRPNWGPNDPRRRSRNLGKVIDTLTCGFAD LMGYIPVVGAPXGGVAXALAHGVXXIEDXVNYATXNLPXXSXSIXLLALLSCLTTPASAAHYTNKS GLYHLTNDCPNSSIVYEAETLILHLPGCVPCVKXXNQSRCWVQASPTLAVPNASTPVTGFRKHVDI MVGAAAFCSAMYVGDLCGGLFLVGQLFTLRPRMHQVVQECNCSIYTGHITGHRMA

SEQ ID NO. 48 (VN12, 7d)

MSTLPKPQRKTKRNTNRRPMDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQARGRRQPIP KVRQNQGRTWAQPGYPWPLYGNEGCGWAGWLLSPRGSRPDWXPNDPRXRSRNLGKVIDTLTCGFAD LMEYIPVVGAPLGGVAAELXHGVRAIEDGINYATGNLPGCSFSIFXLALLSCLTTPASALNYANKS GLYHLTNDCPNSSIVYEANGMILHLPGCVPCVKTGNLTKCWLSASPTLAVQNASVSIRGVREHVDL LVGAAAFCSAMYVGDLCGGLFLVGQLFTFRPRMYEIAQDCNCSIYAGHITGHRMA

SEQ ID NO. 42 (FR1, 9a)

MSTLPKPQRKTKRNTNRRPMDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KVRQPTGRSWGQPGYPWPLYGNEGCGWAGWLLSPRGSRPNWGPNDPRRRSRNLGKVIDTLTXXLAD LMGYIPVLGGPLGGVAAALAHGVRAIEDGVNYATGNLPGCSFSIFLLALLSCLTTPASAIQVKNAS GIYHLTNDCSNNSIVFEAETMILHLPGCVPCIKAGNESRCWLPVSPTLAVPNSSVPIHGFRRHVDL LVGAAAFCSAMYIGDLCGSIILVGQLFTFRPKYHQVTQDCNCSXNXGHVTGHRMA

SEQ ID NO. 50 (NE98, 10a)

 ${\tt MSTLPKPQRKTKRNTNXRPQDVKFPGGGQIVGGVYVLPRRGPQLGVRAVRKTSERSQPRSRRQPIPRARRTEGRSWAQPGYPWPLYGNEGCGWAGWLLSPRGSRPSWGPNDPRRR}$

SEQ ID NO. 52 (NE98, 10a)

DGINFATGNLPGCSFSIFLLALFSCLLTPTAGLEYRNASGLYMVTNDCSNGSIVYEAGDIILHLPG CVPCVRSGNTSRCWIPVSXTVAVKSPCAATASLRTHVDMMVXAATLCSALYVGDLCGALFLXGQGF SWRHRQHWTVQDCNCSI

SEQ ID NO. 54 (BNL1, 1d)

STVTENDIRVEESIYQCCDLAPEARKAIKSLTERLYIGGXLTNSKGQNCGYRRCRASGVLTTSCGNTLTCYLKARAACRAAKLRDCTMLVCGDDLVVICESAGVEEDAANLRA

SEQ ID NO. 56 (BNL2, 1d)

 ${\tt STVTENDIRTEXSIYQCCDLAXEARKAIKSLTERLYVGGPLTNSKGQNCGYRRCRASGVLTTSCGNTLTCYLKARAACRAAKLQDCTMLVCGDDLVVICESAGVEEDAANLRV}$

SEQ ID NO. 58 (FR17, 1d)

STVTENDIRVEESIYQCCDLAPEARKAIKSLTERLYIGGPLTNSKGQNCGYRRCRASGVLTTSCGNTLTCYLKARAACRAAKLQDCTMLVCGDDLVVICESXGVEEDAANLRV

Figure 3 - continued

SEQ ID NO. 60 (CAM1078, 1e)

MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYVLPRRGPRLGVRAARKTSERSQPRGRRQPIP KERRPEGRSWAQPGYPWPLYGNEGCGWAGXLLSPRGSRPSWGPTDPRRRSRNLGKVIDTLTCXFAD LMGYIP

SEQ ID NO. 62 (CAM1078, 1e)

STVTEADIRTEESIYQCCDLHPEARVAIKSLTERLYVGGPLTNSKGENCGYRRCRASGVLTTSCGN TLTCYIKALAACRAAKLQDCTMLVCGDDLVVICESVGTQEDAASLRA

SEQ ID NO. 64 (FR2, 1f)

STVTESDIRTEESIYQCCDLDPEARKAIRSLTERLYIGGPLTNSKGQNCGYRRCRASGVLTTSCGN TLTCYIKARAACRAAKLQDCSMLVCGDDLVVICEIEGXXEDPSXXXX

SEQ ID NO. 66 (FR16,1q)

MSTNPKPQRKTKRNINRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KARRSEGRSWAQPGYPWPLYGNEGMGWAGWLLSPHGSRPSWGPSDPRRRSRNLGKVIDTLTCGFAD LMGYIPLVGAPLGGVARALAQGFRDL

SEQ ID NO. 68 (FR16,1g)

XXVTESDIRVEXSIYQCCDLAPEARVAIKSLTERLYVGGPLTNSKGQNCGYRRCRASGVLTTSCGN TLTCYLKAAAACRAAKLRECTMLVCGDDLVVICESAGVQEDAASXXX

SEQ ID NO. 70 (BNL3, 2e)

STVTERDIXTEESIYQACSLPEQARTAIHSLTERLYVGGPMMNSKGQSCGYRHCRASGVLTTSMGN TITCYIKALAACKAAGIVAPTMLVCGDDLVVISESQGVEEDDRNLXX

SEQ ID NO. 72 (FR4, 2f)

STVTERDIRTEESIYLACSLPEQARTAIHSLTERLYVGGPMMNSKGQSCGYRRCRASGVLTTSMGN TITCYVKALAACKAAGIVAPTMLVCGDDLVVISESQGAEEDERNLRV

SEQ ID NO. 74 (BNL5, 2h)

STVAERDIRTEESIYLACSLPEQARTAIHSLTERLYVGGPMMNSKGQSCGYRRCRASGVLTTSMGN TITCYVKALAACKAAGIVAPTMLVCGDDLVIISESQGTEEDERNLRV

SEQ ID NO. 76 (FR13,2k)

MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLXCRXPRXXXCATXKTXEQSQPRGRRQPIP KDRXTTGKSWGRPGYPWPLYGNEGLGWAGWLLSPRGSRPSWGPTDPRHRSRNLGKVIDTLTXGFXD LMGYIPVVGAPVXGVARALAHGVRVLEDGINYETGNLPGCSFSISLLALLSITXPVSAVEIKNTXN TYMVTNDCSNXSITWQLXXAVLHVPGCVPCEREGNSSRCWIPVTPXVXVSRPGALTEGLRSHIDTI VASATFCSALYIGDVCGAIMIAAQVVIVSPEHHHFVQDCNCSIYPGHITGPRMX

SEQ ID NO. 78 (FR13, 2k)

STVTERDIRVEESVYLSCSLPEEARAAIHSLTERLYVGGPMQNSKGQSCGYRRCRASGVLTTSMGN TLTCYLKAQAACRAAGIVAPTMLVCGDDLVVISESQGTERDENNLRP

Figure 3 - continued

SEO ID NO. 80 (FR18,21)

STVTERDIRNEESIFLACSLPEEARTVIHSLTERLYIGGPMMNSKGQSCGYRRCRASGVFTTSMGN TITCYVKAMAACRAAGIDAPTMLVCGDDLVVISESQGTEEDERNLRV

SEO ID NO. 82 (PAK64,3g)

STVTEQDIRVEEEIYQCCDLEPEARRAIKSLTERLYVGGPMFNSKGLKCGYRRCRASGVLPTSYGN TITCYIKARAAARAGLQDPSFLVCGDDLVVVAESCXVDEEDRAALR

SEO ID NO. 84 (BNL8,4k)

STVTEKDIRPEEEVYQCCDLEPEARKVITALTERLYVGGPMHNSKGDLCGYRRCRASGVYTTSFGN TLTCYLKASAAIRAAGLRDCTMLVCGDDLVVIAESDGVEEDNRALXA

SEO ID NO. 86 (BNL12,41)

STVTEKDIRVEEEIYQCCDLXPEARKAISALTEXLYLGGPMYNSKGELCGYRRCRASGVYTTSFGN TVTCYLKATAATRAAGLKDCTMLVCGDDLVVIAESEGVEEDSQPLRA

SEQ ID NO. 88 (EG81, 4m)

STVTERDIRVEEEVYQCCDLEPEARKAISALTERLYVGGPMFNSKGDLCGYRRCRASGVYTTSFGN TLTCYLKATAATRAAGLKDCTMLVCGDDLVVIAESDGVDEDRRALQA

SEO ID NO. 90 (VN13,7a)

STVTERDVQTEHDIYQCCKLEPAARTAITSLTDRLYXGGPMXNSKGQACGYRRCRASGVLTTILAN TLTCYLKAQAACRAAGLKDFDMLVCGDDLVVISESLGVSEDTSALRA

SEO ID NO. 92 (VN4,7c)

STVTERDIXTEHDIYQCCQLDPVARKAITSLTERLYCXGPMMNSRGQSCGYRRCRASGVLTTSLGN TLTCYLKAQAACRAAKLKNYDMLVCGDDLVVIAESGGVSEDVDALRA

SEO ID NO. 94 (VN12,7d)

SSVTERDIRTEHDIYQCCQLDPVARKAITSLTERLYCGGPMYNSRGQSCGYRRCRASGVFTTSLGN TMTCYLKAQAACRAXKLKNFDMLVCGDDLVVIAESGGVPEDAGALRV

SEQ ID NO. 96 (FR1,9a)

STVTGRDIRTEXDIYLSCQLDPEARKAIKSLTERLYVGGPMYNSKGQLCGQRRCRASGVLPTSMGN TITCFLKATAACRAAGFTDYDMLVCGDDLVVVTESAGVNEDIANLRA

SEO ID NO. 98 (NE98, 10a)

STVTEQDIRVELSIFQACDLKDEARRVITSLTERLYCGGPMFNSKGQHCGYRRCRASGVLPTSFGN TITCYIKAKAATKAAGIKNPSFLVCGDDLVVIAESAGIDEDKSALRA

SEO ID NO. 100 (FR14,11a)

STVTERDIRTEESIYLSCQLPEEARKAIKSLTERLYVGGPMENSKGQACGYRRCRASGVFTTSLGN TMTCYIKAKAACKAAGIVDPVMLVCGDDLVVISESKGVEEDQRDLRV

43/74

SEQ ID NO. 102 (FR15,11a)

STVTERDIRTEESIXXACQLPEEARKAIKSLTERLYVGGPMENSKGQACGYRRCRASGVFTTSLGN TMTCYIKAXAACKXAGIVDPVMLVCGDDLVVISESKGVEEDQRDLXX


SEQ ID NO. 104 (FR19,11a)

 ${\tt MSTNPKPQRQTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRVGVRATRKTSERSQPRGRRQPIP}\\ {\tt KVRRTTGR}$


SEQ ID NO. 106 (FR19,11a)

STVTERDIRTEESXYLACQLPEEARKAIKSLTERLYVGGPMENSKGQACGYRRCRASGVFTTSLGN TMTCYIKAKAACKAAGIVDPVMLVCGDDLVVISESKGVEEDQRDLRX

4. Core/El amino acid alignment	e Type SEQ	מד (MSTNPKPQKKNKRNTNRRPODVKFPGGGOTVGGTVTT BBB CTLL CTLL CTLL CTLL CTLL CTLL CTL	(ZR-TXXXXX	1d 6R-TV	le 10/60R-T	12	1g 66	2a		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	14	18	76	T-T	3h	 44	0 P	4 b + 5 m	10	E	W	4;	4k 28		H-M	461'	F-8I	48L	42LR-TM	LR-TXX	104RQT	
Figure 4.	Isolate	HCV-1	H 777	DICV LO	DIVILI	7TNG	CAM1078	FR2	FR16	HC-J6	HC-J8		BNT3			SH NZI.1		DK13		_		HPCCOREZ	HPCCOREZC	GB724	BNL7	BE95	HK2	VN13	VN4	VN12	FRI	אבים 10 מים	EKIY	




	LSU RGSRPSWGPTDPRRSRNLGKVIDTI,TCGFADT,MGYTP1,VGAP1,CGAAPA		N		XX09	ш−5N	- 1	ΛH	T		11-11-1111	X1X	\$!! !!]			NNN		N		1	AA		NNKKKKK	N	XXXXX	N	;	NNXXIVI-GV-A-A-	N
SEQ	1		۲3	9	0	12	99					14/	18										28			46	44	48		20
Type	1a	1p	1q	1q	1e	1£	1g	رع ه ر	5p	2 0	5g	2e	2£	7, X	3p	4 0	4q	4e	4 e	4 £	4 g	4h	4 k	Бa	6 a	7a		7d	σ (10a
Isolate	HCV1	HCV-J	BNL1	BNL2	CAM1078	FR2	FR16	HC-J6	HC-78												6B249 26		BNL7	BE95	HK2	VN13	VN4	VN12	FR1	NE98



I-SAXULXXII-SAXI		-AII·	[XXXX-X-X-XTAHYI-KS-	IX	-A1I-K-AS-I I-EAS
	į	AI			T W

444 52 52


44x 58a 58a 70 70 98a 10a



```
40
            4 4 4 C
5 C C C
41
5a
5a
7c
7d
9a
BNL12
GB724
BE95
BE100
HK2
VN4
VN12
FR1
NE98
```

757	PATQLRRHIDLLVGSATLCSALYVGDLCGSVFLVGQLFTFIS		VT-AFRM	-Su	O	AA-MILS-A-MVO		AII-MVVAIM-AA-VVVVO	0H-		ATI-MIA-MIAA-VAVV	VIV-A-MIAA-VVIV	FA-MS-F-MT		1	-VAM	-V-MAROAF-A	-VMA	OMG3VVKMV-	VMAMF-IG	S-VMAVIG				MIGI	1	XX	N-WIJI	Λ. Υ. Υ. Ε.	V-87M45H1	X-X-X1VX-X-X-X-X	Z-V-MAMMMMMN-S		S-VMAVI-XGI,	-MGI
SEQ	1	4	ထ	12						16	18	20	24	92	9/																30	32	34	36	38
Type	1a 1b	1q	19 19	1£	2a	2p	N O	2c	7 7 7	2 6	2£	2g	5h	,† 7	2, X	3a	3p	4 a	4a	4p	4°C	4 C	4°C	4q	4e	4 e	4 £	4 £	4 g	4h	4 k	4 k	4 K	4 k	4 k
Isolate	HCV-1 HCV-J	BNL1	BNL2	· FR2	HC-J6	HC-J8	CH610	S83	NE92											21	GB116	≪ GB215	GB358	DK13	CAM600	CB809-2	CAMG22	CAMG27	GB549	GB438	BNL7	BNT8	BNL9	BNL10	BNL11

LSA-LMSVVMASGAMQ VDA-LESFVMAVGAMQ LGAVTAPAV-Y-A-G-AAAL	AST-V-GF-K-V-IMA-AFM
4 ,	4444 522 488
4 4 70 70 70 4 12 12 12 12 12 12 12 12 12 12 12 12 12	7င 7d 9a 10a
GB724 BE95 BE100 HK2	VN4 VN12 FR1 NE98



1	A	aa	A-				
-L	-S	-S	-1	-I	A-	ANX	ı
D	1		\ 			V-7-	,

524 52 52

44 × 5 a 6 a 7 C 7 C 7 C 9 a 1 0 a 1 0 a

GB724 BE95 BE100 HK2 VN4 VN12 FR1

		7981 CTCCACAGTCACTGAGAGCGACATCCGTACGGAGGAGGCAATCTACCAAA	AGATAT	A	GTATGTC	GTBAGGBB		- 1	NA	NNNNNNN	AC	ACGBA-A-AAAAAAAAAA	GAATAA-N	ACAGTAA-A	991 8 -> 1 YY	- AAMGTTATEA-AA-	AA-GA-G	ACAGA-GGTAAGA	ATA-GAGA	ACAGTA-GGTAAA-AAA	
Figure 5. NS5B nucleotide alignment	SEQ TD 7033	_	AG	NACA	3GTB	.5GT	7GTp	1AGAGC	3 NA	NNNNNNN L	AC	ACG			73AG-G			ATACP	TTACAT	81TTACF	
NS5B	Type S	la 1	1b	1b	1d 5	1d 5	1d 5	1e 6	1£ 6	1g 6	2a	2 b					21 7	3a	3b		
Figure 5	Isolate	HCV-1	HCV-J	BE90	BNL1	BNL2	FR17	CAM1078		FR16						_	FR18	$_{ m T1}$	19	PAK64	
										SU	BS	TI	ſŰ	ſΕ	SH	EE	Τ(RL	ILE	26)

```
---T--C-----A----G-----A--G--A--AT-C----T-TG-
                                                         ----A----G--T--A--A--A--AT-C--Y--T-TGG
SEQ
ID
                                   89
91
93
95
97
101
     GB48
GB116
GB215
GB358
GB358
GB809
GB549
BNL12
EG81
CHR18
VN13
VN13
VN13
VN13
FR1
FR1
FR1
FR15
                                    SUBSTITUTE SHEET (RULE 26)
```

Isolate

Isolate

GITGIGACCICGACCCCAAGCCCGCGIGGCCAICAAGICCCICACCGAG ------T-G-C----G----A-GCA-----A-G---G----A -C----T--A-----G-G-T---AA-----A-G-----A --T--G--AG-G--T--GAA---G----GCG-T----A---C----TC-T-ACC-GAG--G---A-AACT----AC-C--AT-G--T--------T-G-C----G-G---A-ACA-----A-----G----A------T-G-C---G-G--T---AA-----A-----G-------A--A--T------GC----G----A--T--A------TT-G--T -C-----A----G-C----G-G--T-----A------G----T C----TC-T-GCC-GAGG-G----A-ACT----AC-C--A--G--T CC--CTC-T-ACC-GAG------AACT----AC-T--AT-G--T CC---TCA--TCC-GAGG-G----A-CT----AC-C--A--T CC--CTCGT-GCC-GAGG-G----GACT-T---AC-T--G---T L-----D--CC--CTC-T-ACC-GAG--G----GACT----AC-T--AT-A--T -C----T-G-C---G-G-----AA-----A-----G----C----TCT--GCCT-AAG----A-AACT-T---AC-C --G--GG-G--TA-ACG---A------T-G-C---YG-G-----AA---69 77 77 79 Type CAM1078 HC-J6 HC-J8 BNL3 HCV-1 HCV-J PAK64 FR2 FR16 BNL5 FR13 FR18 **BE90** BNL2 FR17 BNL1 FR4 SUBSTITUTE SHEET (RULE 26)

```
8031
                      CA-TGT--T-GC-G--TG-G--G--T----A-ACG--A---A--C-A
-C---A-GT-G--G---GC--A--GACA------CA--G-T--T--C
-C---CC-A--T-----GGTG--A--AA--T--T-CA--T--G--T---
                                                                   -C--CC-AT-A--T--GGT---A--GAAA----T-CA--T
SEQ
ID
                                                       889
931
957
101
103
                                    83
85
87
_{\mathrm{Type}}
          Isolate
        GB48
GB116
GB215
GB358
GB809
GB549
BNL12
EG81
CCHR18
VN13
VN13
VN13
VN13
FR1
FR1
FR1
FR1
                                                          SUBSTITUTE SHEET (RULE 26)
```

8081 AGGCTTTATGTTGGGGGCCCTCTTACCAATTCAAGGGGGGAGAACTGCGG ----A-GTT---CAGC-A----CCC-A--T----A--C--CA-A--C--G--GA-G-TG--CAGC-AA--CC--TC---T-C----G--CA-C--A--T--CA-GTA---CAGT-A----CTCC-G-------G--C--G--A--T--CA-GCAG--CAGC-A----C-ATC----C--G--A--G--CA-G-TG--CAGC-AA---C--TC---C--A--A--G--CA-G-TG--CAGC-A---AC--TC---C-----C----C----T--C--G--T-----G-A----C------A----C--G--A--G--CA-GTT----CAGC-A---CC---C --A--C--C--A--A--G--CA-G-TG--CAGC-AA---C-ATC---G--CA-G--A--CAGC-AA---C-ATC------C----C----AA--AC------G--T----A----C-----A----C---AA---C-------T--C--G--A--C----AA---C-----C--A-----A----C--A--A--C----G--CA-C-------G---A-C -----Y-C 8032 SEQ ID 53 57 61 63 81 $_{\mathrm{Type}}$ BNL2 FR17 CAM1078 Isolate HCV-1 HCV-J HC-J6 PAK64 HC-JB FR2 FR16 BNL5 FR13 FR18 BE90 BNL3 BNL1 FR4 SUBSTITUTE SHEET (RULE 26)

Isolate

```
8081
                                                                                        C-C--G--CTG---A----CA-GTAT--CAGC-A----C-AC-A--T---C-AT-G--CTNC--T--T---CA-GTNT---C--T-AA--TC--GCA--T--
                                                                                                                 C----G--CTGC--C-----CA-GTA---C--TC-A--TC--TCA--T--
                                                                                                        C----G--CIGC--W--G--CA-G-IG--C--CC-I--IC-AICA--I-
                                                        .C--G--C--T--CA-GTA---C--C-A------CCTA-----
-C--G--C-----CA-GCA---CAGC-A---A--CCTT--T--
-CT-G--C-----CA-GTAT--CAGC-AA------CT----
                                          --CA-GCAT--CAGC-A---A--CCTG--T-
                                                                                 C----G--C--T--CA-GITT--CAGC-A---A--CCTA--T-
                                                  ----CA-GCAT--CAGC-A---A--CCTT--
                                                                                                                                 C-----CTG---T--T--A-GTT---CAGC-A---AC-AC---
                                                                          C--CI-G--
                                                --A--C--(
                                  --A--C-
                                         --A--C-
                                                                         --R--(
                                                                  --A--
                                                                                 --A--
 SEQ
ID
                                                                                                89
91
93
97
101
105
                                                                 83
85
87
Type
                 GB48
GB116
GB215
GB358
GB809
GB549
BNL8
BNL12
EG81
CHR18
VN13
                                                                                                               VN12
FR1
NE98
                                                                                                                 SUBSTITUTE SHEET (RULE 26)
```

```
CTATCGCAGGTGCCGCGGGGGGGTACTGACAACTAGCTGTGGTAACA
                                                               -G--T-
                                                         TT-C--C--ATG--G--T
                       --C--TC----T----T
                                                       ---CA-GC-T-----A----T
                   -----R----R-----
                                         ---C---C-A----T--A--
                                                                ----L----C-----
                                                  G--CA-GC-T----C
                                      -----A----T
SEQ
ID
                       53
57
61
63
                                                           69
71
77
79
Type
         CAM1078
FR2
FR16
HC-J6
HC-J8
BNL3
Isolate
        HCV-1
HCV-J
                                                                                          PAK64
                      BNL1
BNL2
FR17
                 BE90
                                                                   BNL5
FR13
FR18
T1
```

SUBSTITUTE SHEET (RULE 26)

```
SEQ
ID
                       89
91
93
97
99
101
105
               83
85
87
Type
    Isolate
   GB48
GB116
GB215
GB358
GB809
GB549
BNL12
EG81
CCHR18
VN13
VN13
VN12
FR1
FR1
FR1
FR1
FR15
                           SUBSTITUTE SHEET (RULE 26)
```

8132 CCCTCACTTGCTACATCAAGGCCGGGCAGCCTGTCGAGCCCGCAGGCCTC
SE ID 1D 63 77 77 81
туреран прад прад прад прад прад прад прад прад
I solate HCV-1 HCV-1 HCV-1 HCV-1 BNL1 BNL1 ER17 CAM1078 FR2 FR16 HC-J6 HC-J7 HC-J6 H

8132 -AGGGGGGG -AGGGGGG -AGG
SEQ ID 101 101 105
T 4444444447777777777777777777777777777
Isolate GB48 GB116 GB215 GB358 GB809 GB809 GB809 GB809 GB809 GB816 GB115 GB116 GB115 GB115 GB115 FR1 FR1 FR1 FR1 FR1 FR15 FR19

```
8231
             CAGGACTGCACCATGCTGTGTGGCGACGACTTAGTCGTTATCTGTGA
                                                                            ----G--I--C---CA--
                                                                                                                                       --C--A--T--G--G--AG-GGC--
                                                                                      GT--C-CC----T----G-----T----C----T-
                     ---C--G--L--C-L--
                                                                                GT----CCTGTT---T-G-----A----C
                                                                          ATT-CGCC---A----G--A--C----T
                                                 -----C--A--
                                                                                                           GTT-CACC---A----G-----
                                                                                             GTT-C-CC---G----G----C
                                                                                                    GTT-CTCC---G----G--T--
                                                                                                                  G-C-CC---A---T-G
                                                                                                                                       --A---CCAT-AT-C--T
                            ----9----
                                                                   G---A----A---
                                                             ----T----
SEQ
ID
                                 53
57
61
63
                                                                                       69
71
77
79
                                                                                                                                      81
Type
             33212766
33212766
33212766
33212766
33212766
33212766
33212766
            HCV-1
HCV-J
BE90
BNL1
BNL2
FR17
CAM1078
FR2
FR2
FR2
HC-J6
Isolate
                                                                                                                                     PAK64
                                                                                            FR4
BNL5
FR13
FR18
T1
                                                                                          SUBSTITUTE SHEET (RULE 26)
```

```
8231
                  A-AA---ATGA---T-A--C
                                        --9--D-D-----
                                                      ACA--T-ATGA----T-G
A-AA-TCCAT-AT-C--T
       AGA----
AGA----
AGA----
SEQ
ID
                             83
85
87
Type
       Isolate
       GB48
GB116
GB215
GB358
GB809
GB549
BNL12
EG81
CCHR18
VN13
VN13
VN12
FR1
FR1
FR1
FR15
                                                SUBSTITUTE SHEET (RULE 26)
```

```
AAGCGCGGGGGTCCAGGAGGACGCGGCGAGCCTGAGAGCC
G--T----AAC-----T----GC---AC----
                                                                  ----AGA--AGCT---C---
                                                                     ---G-ATAG-GCAGC
                                                                    G--TIGC-KC--IG-I--
SEQ
ID
                 53
55
61
63
63
                                            69
71
73
79
Type
       BNL1
BNL2
FR17
CAM1078
FR2
FR16
HC-J6
HC-J8
BNL3
FR4
BNL5
FR13
FR18
Isolate
         HCV-J
BE90
                                                                   PAK64
      HCV-1
                                             SUBSTITUTE SHEET (RULE 26)
```

```
8271
                   G--T-GA--A---TCT----T-TT-ACGC----C---A
G----GA--A----CT----T--C-G-GC----C---T-
                                                                                    ---AA-AGCGC-T----T
                                                                              --TATC--T-A----C----
                                                                                                    -CAACGAGA---AC--NT-
                                                                                          --CA-CG-GA---AC---T
            --AAACGACC---CG---
                 --AAACGAGC---CG---
                                                                                              --CA-CGAGA---AC
                                                                              --T--A---A-C--
                 ----AT--C-AG---
           G---AT--C--AG--
      8232
 SEQ
ID
                                                             89
91
93
93
101
101
                                         83
85
87
Type
           1110a
1110a
1110a
1110a
1110a
Isolate
          GB48
GB116
GB215
GB358
GB809
GB549
BNL8
BNL8
BNL12
EG81
CHR18
                                                                       SUBSTITUTE SHEET (RULE 26)
                                                                  VN4
```

NS5B amino acid alignment	2,70	2694 STVTESDIBTER IVOCORI DRONDEN TWO THE STREET	N		! - 	XX		1							TO-XXXXXXXXXXX	! ! ! !		RVSV-1,S-S-DEEA	i	; !	CK_AQ		HEE-E-EKSDTMVV	QVEE-E-R
B amin	SEQ	ا ن			54	56	58	62	64	89					70	72	74	78	80					82
- 1	Type	1a	1p	1c	1d	1d	1d	le	1£	1g	6 2	5 p	2	2d	2e	2£	Sh	2k	21	3a	3а	3a	3b	3g
Figure 6.	Isolate	HCV-1	HCV-J	2TY4	BNL1	BNL2	FR17	CAM1078	FR2	FR16	HC-J6	HC-JB	C ARG8	S NE92	ENT3	II FR4	S BNL5	H FR13	FR18	3. BR34	108 BR 36	H BR33		PAK64

	47-11
888 89999111 88 00011 0001 0001	
2 4 4 4 C C C C C C C C C C C C C C C C	
GB48 GB116 GB116 GB215 GB328 GB328 GB349 GB438 CAR4/12 CAR1/5(CAR1/5(CAR1/5(CAR1/8 BNL12 BNL13 BNL12 BNL12 BNL12 BNL13 BNL12 BNL13 BNL12 BNL13 BNL12 BNL13 BNL12 BNL13 BNL12 BNL13 BNL13 BNL12 BNL13 BNL1	E 26)

2695	ZOOS YRRCRASGVI, TTSCGNTI, TCV TKADAACDAACT ORGENTINGERS	THE THE TENER OF T			\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\			4			, ,	V			VTX			<u> </u>		- 1 - 4 - 4		K-DSE	- !	
Type SEQ)			54	56	28	62	64	89					70	72	74	78	80					82	
Type	la	1p	1c	1d	1d	1d	1e	1£	1q	(2) (a)	5 p	2C	5d	3e	Z£	5h	2k	21	3a	3a	3a	36	39	
isolate	HCV-1	HCV-J	2TY4	BNL1	BNL2	FR11	CAM1078	FR2	FR16	HC-J6	HC-J8	ARG8									BR33		F PAK64	6)

```
90
92
94
98
100
102
          84
86
88
SUBSTITUTE SHEET (RULE 26)
```

```
-Q-TE--ERN---
-Q-NE--ERN---
-Q-TE--ERN---
-Q--E--DRN-
-Q-AE--ERN--V
                      ----E---N---
----E---N--V
-X--E----N--V
-V-T-----
                                                                         -Q-TE--ERN--V
-Q-TER-ENN--P
-Q-TE--ERN--V
    2745
SAGVQEDAASLRA
                                                                                                    -C--E--R-A---
             ---T----A---
                  V----T---
                                         IE-XX--PS
SEQ
ID
                                                                70
72
74
78
80
                       554
62
68
68
Type
         HCV-1
HCV-J
BE90
BNL1
BNL2
FR17
CAM1078
FR2
FR2
HC-J6
HC-J6
HC-J8
NE92
BNL3
FR4
BNL3
FR4
BNL3
FR13
FR18
BNL3
FR13
FR18
Isolate
                                                                SUBSTITUTE SHEET (RULE 26)
```

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
FADED TEXT OR DRAWING
BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
GRAY SCALE DOCUMENTS
LINES OR MARKS ON ORIGINAL DOCUMENT
REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
OTHER:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

